



2010

Aflatoxin in raw peanut kernels marketed in Malaysia

Follow this and additional works at: <https://www.jfda-online.com/journal>

Recommended Citation

Arzandeh, S.; Selamat, J.; and Lioe, H. (2010) "Aflatoxin in raw peanut kernels marketed in Malaysia," *Journal of Food and Drug Analysis*: Vol. 18 : Iss. 1 , Article 5.
Available at: <https://doi.org/10.38212/2224-6614.2222>

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

Aflatoxin in Raw Peanut Kernels Marketed in Malaysia

SAHAR ARZANDEH¹, JINAP SELAMAT^{1*} AND HANIFAH LIOE^{1,2}

¹ Centre of Excellence for Food Safety Research (CEFSR), Faculty of Food Science and Technology,

Universiti Putra Malaysia, UPM 43400, Serdang, Selangor, Malaysia

² Department of Food Science and Technology, Faculty of Agricultural Technology, Bogor Agricultural University, Bogor 16002, Indonesia

(Received: June 26, 2009; Accepted: January 16, 2009)

ABSTRACT

The occurrence of aflatoxin in eighty-four samples of raw peanut kernels which are randomly collected from Malaysian supermarkets was examined. Analysis for aflatoxin was performed by solvent extraction and immunoaffinity clean-up followed by the determination using high performance liquid chromatography equipped with post-column photochemical reactor for enhanced detection and fluorescence detector. A detection limit of 0.01-0.09 ng/mL and a quantification limit of 0.04-0.30 ng/mL were obtained. The aflatoxin concentrations ranged from not detected to 97.28 ng/g in all samples investigated. About 78.57% of the samples were contaminated with aflatoxin, of which 10.71% exceeded the maximum tolerable limit of 15 ng/g set by the Codex. Average recoveries of the aflatoxin analysis were acceptable which were in the range of $74.85 \pm 8.83\%$ for AFG₂ at the concentration of 0.15 ng/mL and $103.91 \pm 6.45\%$ for AFB₂ at the concentration of 0.15 ng/mL. The average daily intake estimated for total aflatoxins was 10.69 ng/kg body weight. There was a significant difference ($P < 0.05$) in aflatoxin content between brands and locations.

Key words: Aflatoxin, peanut, food safety, HPLC with fluorescence detection, immunoaffinity clean-up

INTRODUCTION

Aflatoxins are secondary metabolites produced by many strains of *Aspergillus flavus*, *Aspergillus parasiticus* and the rare *Aspergillus nominus*⁽¹⁾. After the death of 100,000 turkeys in the UK in 1960s owing to toxic metabolites coming from fungi⁽²⁾, aflatoxins have been a major concern as a carcinogenic, mutagenic and immunosuppressive agent of feeds and foods⁽³⁾. Various agricultural commodities including peanut, corn, cottonseed, Brazil nut, pistachio nut, fig, spice and copra are likely to be contaminated by aflatoxins⁽⁴⁾. In comparison to the other agricultural commodities, peanut is very susceptible to aflatoxins contamination. It could be due to the kernels develop and mature beneath the surface, and domination of *Asp. flavus* of the peanut field soil⁽⁵⁾.

Despite the fact that 20 aflatoxins have been identified, only 4 of them, the aflatoxin B₁, B₂, G₁ and G₂ (AFB₁, AFB₂, AFG₁ and AFG₂), are fundamental contaminants of a diversity of foods and feeds⁽¹⁾. AFB₁ has been classified by the International Agency for Research on Cancer (IARC) as a group 1 carcinogen,

mainly to cause liver cancer⁽¹⁾, whereas AFB₂, AFG₁, AFG₂ are classified as possible carcinogens to humans⁽⁶⁾. Since aflatoxins are potent source of health hazards to both human and animals and they are causing lots of economic losses, attempts have been made to study the aflatoxins occurrence in many parts of world and to completely annihilate the toxin or diminish its content in foods and feedstuffs⁽⁷⁾.

The two probable reasons for aflatoxin contamination in peanut are severe late-season drought stress happening in the field (pre-harvest) and the existence of undesirable moisture and temperature conditions during storage (post-harvest)⁽⁸⁾. Peanut shell penetration by molds is facilitated by physical damage; hence, aflatoxins contamination will occur⁽⁹⁾. Malaysia, a tropical country with an average temperature of 28 to 31°C and heavy rainfall throughout the year, is appropriate for fungal growth. However, in the dry season the relative humidity is 50 to 60%, in contrast to 70 to 80% in wet seasons; hence mold growth and aflatoxins production in products such as peanut stored under these conditions will increase⁽¹⁰⁾.

Therefore, the purpose of this study was to examine the occurrence and concentration of aflatoxins in packaged plastic bag raw peanut kernels marketed in

* Author for correspondence. Tel: +60-3-8946-8393;

Fax: +60-3-8942-3552;

E-mail: jinap@food.upm.edu.my; sjinap@gmail.com

Malaysian supermarkets located in four different areas using an immunoaffinity column AflaTest_{WB} (IAC) clean-up and high performance liquid chromatography (HPLC) with fluorescence detector and to estimate the daily intake of this toxicant from peanut consumption.

MATERIALS AND METHODS

Mixed aflatoxin standards with AFB₁ and AFG₁ concentration of 300 ng/mL, AFB₂ and AFG₂ concentration of 1000 ng/mL, individual AFB₁ with 20000 ng/mL, and AFB₂, AFG₁ and AFG₂ with concentration of 3000 ng/mL were purchased from Supelco (Bellefonte, PA, USA). All solvents used for the experiments were of HPLC grade and supplied by Merck (Darmstadt, Germany). AflaTest_{WB} immunoaffinity columns (IAC) with 1 mL volume were purchased from Vicam Company (Watertown, MA, USA).

I. Sampling

A total of 84 samples of packaged plastic bag raw peanut kernels were purchased randomly from different supermarkets in four locations (north, south, east and west region) of state of Selangor, Malaysia from April to August 2008. A wide range of brands were covered to ensure that the survey was representative of the range of products available to consumers in Malaysia. About 4 kg of samples were randomly collected from each supermarket. The samples were thoroughly mixed and went through quarter sampling to make a representative sample (100 g); the representative samples were then immediately transferred to dry clean polyethylene bags and stored at -18°C prior to analysis.

II. Extraction and Clean up

Mycotoxins should be separated from the solid phase of the matrix and distributed into the liquid phase⁽¹¹⁾. Aflatoxins were extracted and determined using the AOAC official method 991.31⁽¹²⁾ with minor modification. The representative sample (100g) was ground using a Waring blender (Vicom, Watertown, MA, USA) for about 3 min. Twenty-five grams of ground peanut samples and 5 g of NaCl were blended with 125 mL of methanol/water (70:30, v/v) for 2 min. Fifteen milliliters of the extract was diluted with 30 mL water after being filtered on a 24-cm fluted filter paper (Vicom, USA). Finally it was filtered on a Whatman glass microfiber filter (934-AH, Maidstone, UK). Fifteen milliliters of the filtrate was applied to the IAC containing monoclonal antibody specific for AFB₁, AFB₂, AFG₁, and AFG₂ to be purified. The column was washed with 20 mL of deionized water, then aflatoxins were eluted from the column with 1.0 mL of methanol and the eluted fraction was diluted twice with deionized

water, and then stored in a vial at -18 to -20°C.

Aflatoxins extract in the methanol-water solution were determined by HPLC method with fluorescence detector after using a post-column photochemical reactor for enhanced detection (PHRED) (Aura Industries, N.Y, USA).

III. HPLC Determination of Aflatoxins

A HPLC method was used for aflatoxin analysis of all samples, using a reverse phase symmetry C18 column (Merck, Darmstadt, Germany) with a dimension of 25 cm × 4.6 mm, and 5 μm particle size, running on a Waters 2475 HPLC equipped with a fluorescence detector operated at an excitation wavelength of 365 nm and an emission wavelength of 435 nm. The mobile phase was a mixture of water/methanol/acetonitrile (54:29:17, v/v/v) with a flow rate of 1 mL/min. A post-column PHRED was used to enhance the natural fluorescence of AFB₁ and AFG₁ and to improve detection⁽¹³⁾. The PHRED was located between the LC column and the detector; encompass a lamp holder, a 254 nm low-pressure mercury lamp, and a holder for the knitted reactor coils. During photolysis, AFB₁ and AFG₁ are converted to hemiacetals meaning AFB_{2a} and AFG_{2a}, respectively. Due to the very low detectability of AFB₁ and AFG₁, the post column derivatization is used to make them detectable by the detector. The reactor coils are made of polytetrafluoroethylene (PTFE) that is transparent to the 254 nm UV light with 25m length⁽¹⁴⁾.

For aflatoxin identification, linearity, accuracy, repeatability (RSD_r), reproducibility (RSD_R), limit of detection (LOD) and limit of quantification (LOQ), linear equation and coefficient of regression (R²) of the analytical method were determined. Linearity was estimated by injecting triplicate aflatoxin standards. Recovery studies were carried out by spiking aflatoxins in three replicated peanut samples at concentrations of 0.50, 5.00, 30.00 ng/mL and 0.15, 1.50, 9.00 ng/mL of AFB₁, AFG₁ and AFB₂, AFG₂ respectively. The limit of detection (LOD) and the limit of quantification (LOQ) were estimated using the 3xstandard deviation and 10xstandard deviation, respectively, calculated by 7 times injection of standards having the lowest concentration to be detected into the HPLC.

IV. Estimated Daily Intake

The estimated daily intake (EDI) values of aflatoxin by an adult (ng/kg body weight) were calculated using the average value by each type of peanut, i.e. EDI in ng/kg body weight = mean concentration of aflatoxin (ng/g) multiplied by the amount of peanut consumed/day (g) and divided by the average weight of an individual (60 kg)⁽¹⁵⁾.

V. Statistical Analysis

The descriptive statistic (mean, standard deviation

and range) and analysis of variance (ANOVA) were employed using Minitab (Version 14, PA., State College, USA). A probability value of 0.05 was used to determine the statistical significance.

RESULTS AND DISCUSSION

I. Method Performance

For aflatoxin analysis, linearity was estimated by injecting triplicate aflatoxin standard solutions at concentrations of 2, 4, 6, 8, 10, 25, 50 and 100 ng/mL for AFB₁ and AFG₁, and 0.6, 1.2, 1.8, 2.4, 3.0, 7.5, 15.0 and 30.0 ng/mL for AFB₂ and AFG₂, respectively. Eight-point calibration curve was built for each individual aflatoxin used for quantification of aflatoxin in peanut samples. The correlation coefficient was more than 0.993 ($R^2 >$

0.993). The recoveries of aflatoxins in peanut samples are summarized in Table 1. The recovery for AFB₁ ranged from 95.53 ± 11.60 to $102.39 \pm 7.64\%$, AFB₂ ranged from 88.73 ± 10.93 to $103.91 \pm 6.45\%$, AFG₁ ranged from 91.72 ± 5.56 to $97.23 \pm 10.96\%$, and AFG₂ ranged from 74.85 ± 8.83 to $78.00 \pm 11.31\%$. The recoveries obtained for aflatoxins were in line with the legislated levels for aflatoxin determination methods described by commission regulation⁽¹⁶⁾.

The chromatogram of the spiked peanut samples has well-separated peaks, as shown in Figure 1(a). The limit of detection (LOD) and limit of quantification (LOQ) are shown in Table 2. The LOD were found to be 0.03, 0.01, 0.09, and 0.06 ng/mL and the LOQ were 0.10, 0.04, 0.30 and 0.20 ng/mL for AFB₁, AFB₂, AFG₁ and AFG₂, respectively. Moreover, the repeatability (RSD_r) obtained for AFB₁, AFB₂, AFG₁ and AFG₂ were 1.62, 4.45, 5.10 and 9.91%, respectively. The reproducibility (RSD_R) found to be 7.92, 7.25, 9.62, and 10.94% for AFB₁, AFB₂, AFG₁ and AFG₂ respectively.

Table 1. Recoveries and relative standard deviations (%) of aflatoxin in spiked samples

Aflatoxins	Concentration of spiked aflatoxin (ng/mL)	Mean recovery ^a ± RSD (%)
AFB ₁	0.50	100.03 ± 6.80
	5.00	102.39 ± 7.64
	30.00	95.53 ± 11.60
AFB ₂	0.15	103.91 ± 6.45
	1.50	96.80 ± 3.91
	9.00	88.73 ± 10.93
AFG ₁	0.50	91.72 ± 5.56
	5.00	92.74 ± 9.22
	30.00	97.23 ± 10.96
AFG ₂	0.15	74.85 ± 8.83
	1.50	77.73 ± 10.50
	9.00	78.00 ± 11.31

^a Mean recoveries were ascertained by assessing three replicate samples at each spiked level.

II. Aflatoxin Occurrence in Analyzed Samples

The concentrations of aflatoxins in all of the peanut




Figure 1. HPLC fluorescence chromatogram of spiked peanut samples and blank sample. (a) Spiked peanut samples with 2 ng/mL of AFB₁ and AFG₁, and 0.6 ng/mL of AFB₂ and AFG₂. (b) Blank sample.

Table 2. Linear equation, LOD and LOQ obtained for quantification of aflatoxin

Aflatoxins	LOD ^a (ng/mL)	LOQ ^b (ng/mL)	Calibration curve	R ²
AFB ₁	0.03	0.10	$y = 8.97x - 11.65$	0.9948
AFB ₂	0.01	0.04	$y = 20.99x - 8.33$	0.9949
AFG ₁	0.09	0.30	$y = 3.18x - 4.33$	0.9951
AFG ₂	0.06	0.20	$y = 6.88x + 0.18$	0.9962

^alimit of detection.

^blimit of quantification.

samples used in the study are shown in Table 3. The results revealed wide variation in aflatoxins concentrations among the 84 samples analyzed. The study found that 78.57% of the samples were contaminated with total aflatoxins concentrations varying from 2.76 to 97.28 ng/g, whereas 10.71% of the samples exceeded the maximum tolerable limit (15 ng/g) set for total aflatoxins in nuts by the Codex⁽¹⁷⁾. It was interesting to note that only 10.71% of the samples were contaminated with AFG₁ and all samples were free from AFG₂. This might be due to the invasion of peanuts by *Asp. flavus* rather than *Asp. parasiticus*⁽¹⁾.

As shown in Table 4, 75, 67.85, and 10.71% of

samples found to be contaminated with AFB₁, AFB₂, and AFG₁ with the mean concentration of 9.00 ng/g, 1.91 ng/g, and 0.38 ng/g respectively.

Samples of brand 1 were found to contain the highest level of aflatoxins contamination (total 48.95 ng/g) and brand 6 contained the lowest (2.89 ng/g). Moreover, the highest contamination of total aflatoxins was found in the south of Selangor state (19.71 ng/g) and the lowest was in the east (4.10 ng/g). This quite high contamination in the south might be due to the bad condition of its storage. The statistical analysis indicated significant difference ($p < 0.05$) between total aflatoxins, location and brands. The statistical differences are shown

Table 3. Concentrations of aflatoxins in 84 peanut samples analyzed by HPLC

Brands	Location	Sample number	AFB ₁ (ng/g) (mean ± SD)	AFB ₂ (ng/g) (mean ± SD)	AFG ₁ (ng/g) (mean ± SD)	AFG ₂ (ng/g) (mean ± SD)	Total aflatoxins (ng/g) (mean ± SD)
B1	1	3	60.67 ± 4.46 ^a	8.75 ± 0.63 ^a	3.52 ± 0.47 ^a	ND	72.94 ± 5.39 ^a
B2		3	2.95 ± 0.11 ^b	1.45 ± 0.59 ^b	ND	ND	4.40 ± 0.68 ^b
B3		3	3.04 ± 0.27 ^b	1.21 ± 0.45 ^b	3.31 ± 0.40 ^a	ND	7.56 ± 1.12 ^b
B4		3	7.93 ± 0.50 ^c	1.58 ± 0.54 ^b	ND	ND	9.51 ± 0.51 ^b
B5		3	ND	ND	ND	ND	ND
B6		3	2.87 ± 0.10 ^b	ND	ND	ND	2.87 ± 0.10 ^b
B7		3	ND	3.47 ± 0.50 ^c	ND	ND	3.47 ± 0.50 ^b
B1	2	3	87.02 ± 4.63 ^a	10.26 ± 0.80 ^a	ND	ND	97.28 ± 5.43 ^a
B2		3	2.76 ± 0.06 ^b	ND	ND	ND	2.76 ± 0.06 ^b
B3		3	2.89 ± 0.21 ^b	ND	ND	ND	2.89 ± 0.21 ^b
B4		3	4.81 ± 0.37 ^b	2.86 ± 0.76 ^c	ND	ND	7.67 ± 1.08 ^c
B5		3	6.46 ± 0.29 ^b	2.69 ± 1.34 ^c	ND	ND	9.15 ± 1.62 ^c
B6		3	6.24 ± 0.56 ^b	2.46 ± 0.74 ^c	ND	ND	8.70 ± 0.34 ^c
B7		3	7.72 ± 0.59 ^b	1.81 ± 0.23 ^c	ND	ND	9.53 ± 0.40 ^c
B1	3	3	2.86 ± 0.15 ^a	0.95 ± 0.11 ^a	ND	ND	3.81 ± 0.25 ^a
B2		3	2.72 ± 0.08 ^a	0.98 ± 0.15 ^a	ND	ND	3.70 ± 0.24 ^a
B3		3	7.84 ± 0.41 ^b	2.41 ± 0.52 ^b	ND	ND	10.25 ± 0.92 ^b
B4		3	8.27 ± 0.66 ^b	2.71 ± 0.65 ^b	ND	ND	10.98 ± 0.48 ^b
B5		3	ND	ND	ND	ND	ND
B6		3	ND	ND	ND	ND	ND
B7		3	ND	ND	ND	ND	ND
B1	4	3	17.81 ± 0.97 ^a	3.98 ± 0.52 ^a	ND	ND	21.79 ± 1.45 ^a
B2		3	2.80 ± 0.10 ^b	0.98 ± 0.13 ^b	ND	ND	3.78 ± 0.22 ^b
B3		3	4.17 ± 0.33 ^c	1.88 ± 0.27 ^c	3.69 ± 0.66	ND	9.75 ± 1.25 ^c
B4		3	2.92 ± 0.20 ^b	0.99 ± 0.15 ^b	ND	ND	3.91 ± 0.32 ^b
B5		3	7.14 ± 0.67 ^d	1.94 ± 0.33 ^c	ND	ND	9.07 ± 0.99 ^c
B6		3	ND	ND	ND	ND	ND
B7		3	ND	ND	ND	ND	ND

1: North

2: South

3: East

4: West

ND: Not Detected

^{a,b,c,d}Similar letters in each column show insignificant differences.

Table 4. Prevalence of aflatoxins in 84 samples analyzed by HPLC

Aflatoxins	Positive samples (%)	Mean (ng/g)	Max (ng/g)	Median (ng/g)	Range
AFB ₁	75.00	9.00	92.07	2.99	0-92.07
AFB ₂	67.85	1.91	11.16	1.12	0-11.16
AFG ₁	10.71	0.38	4.36	0	0-4.36
AFG ₂	0	ND	ND	ND	ND
Total	78.57	11.28	103.23	4.18	ND

ND: Not detected.

in Table 3. The contamination which was detected in the samples is likely due to either the undesirable pre-harvest or adverse storage conditions⁽⁸⁾.

The occurrence of aflatoxin in peanut has been reported by several authors from different countries. In the Philippines, a survey on peanut-based products showed 60% of the samples were contaminated by AFB₁ and AFG₁ in the ranges of 1-244 ng/g and 6-68 ng/g, respectively⁽¹⁸⁾. Juan *et al.*⁽¹⁾ reported a 0.3 ng/g aflatoxin contamination in peanut in the Rabat-Sale area, Morocco in 2008. Haydar *et al.*⁽¹⁹⁾ analyzed the aflatoxin concentrations in Syrian foods and showed 2.7 ng/g AFB₁ contamination in 28.5% of raw shelled peanut. Farombi⁽²⁰⁾ reported 43-1099 ng/g aflatoxin in Brazilian peanuts in 1998, whereas Abdulkadar *et al.*⁽²¹⁾ did not find any contamination in peanuts in Qatar in 2000. Chun *et al.*⁽⁶⁾ showed 0.2 ng/g AFB₁ contamination in 25% of raw peanuts in South Korea and 20-200 ng/g in peanut samples, all from Argentina and Senegal. Due to the availability of peanut in all retail markets and supermarkets throughout Malaysia and its use in a variety of popular Malaysian foods such as satay (meat or chicken with peanut sauce) and rempeyek (traditional cracker), Malaysian people are at risk from the undesirable effects of aflatoxin on their health^(22,23).

The differences of aflatoxin occurrence in different countries could be related to their different weather conditions, and pre-harvest and post-harvest practices; the phenomena is in agreement to what Akbas and Ozdemir mentioned that geographic location, agricultural practices and susceptibility of the products of fungal growth during harvest, storage and processing affect the occurrence of aflatoxin⁽³⁾.

The recoveries from the spiked samples from the current study were different from other studies. Abdulkadar *et al.*⁽²¹⁾ (2000) determined the amount of aflatoxin in different nuts, using HPLC and precolumn derivatization (trifluoroacetic acid), based on AOAC 990.33. The mobile phase was methanol/water/acetonitrile (13:74:13, v/v/v) at the flow-rate of 0.50 mL/min. The average recoveries for pistachio were 87, 95, 93 and 89% and the repeatability values (RSDr) were 6.12, 10.93, 6.97 and 8.31% for AFB₁, AFB₂, AFG₁ and AFG₂ respectively.

The study reported a limit of detection of 0.1 ng/mL.

In Chun *et al.*⁽⁶⁾ (2007) study, nine types of samples were analyzed using HPLC and precolumn derivatization (trifluoroacetic acid), based on AOAC method 990.33. The mobile phase was water/acetonitrile (3:1, v/v) at the flow-rate of 1 mL/min. The recoveries in peanut butter and walnut were 102, 84.8, 102.1 and 83.4% for AFB₁ and AFB₂ at spiking level of 20 ng/mL and AFG₁ and AFG₂ at spiking level of 10 ng/mL, respectively. The precision as determined by a multiple analysis of spiked samples was 7.11, 22.59, 5.42, and 27.75% for AFB₁, AFB₂, AFG₁ and AFG₂ respectively. The range of limit of detection was 0.08-1.25 ng/mL whereas for the limit of quantification it was 0.15-2.50 ng/mL.

Juan *et al.*⁽¹⁾ (2008) studied the occurrence of aflatoxins in dried fruits and nuts using HPLC and precolumn derivatization (trifluoroacetic acid). The mobile phase used was methanol/water/acetonitrile (17:54:29, v/v/v) with the flow rate of 0.8 mL/min. The recoveries were 83.6, 87.3, 88.5, and 89.5% for AFB₁, AFB₂, AFG₁ and AFG₂ respectively. The limit of detection was 0.006 ng/mL for AFB₁ and AFG₁ and 0.015 ng/mL for AFB₂ and AFG₂ whereas the limit of quantification was 0.02 ng/mL for AFB₁ and AFG₁ and 0.05 ng/mL for AFB₂ and AFG₂.

The differences in the recoveries value were possibly due to either interference of the fluorescence properties of the sample matrix in the detection process of this toxin component by HPLC⁽¹³⁾ or different methods of extraction. The results gained in this study indicate that the HPLC method adopted in this research was acceptable.

III. An Assessment of Aflatoxin Exposure in Humans

The amount of peanut consumption may vary considerably from one individual to another. The daily intake of compounds from food consumption is dependent on the compound concentration in food and the amount of food consumed. Results of an official survey have shown that the average Malaysian consumes 56.90 g/day of peanut and the demand for peanut consumption is increasing over the years⁽²⁴⁾. Based on this input

and the mean concentrations of total aflatoxins in peanut found in this study (11.28 ng/g), for 60 kg as the average of body weight, the ingestion of total aflatoxins was 10.69 ng/kg body weight per day.

CONCLUSIONS

This study showed as many as 78.57% of the 84 peanut samples were contaminated with aflatoxins, of which 10.71% of the samples exceeded the maximum tolerable limit for total aflatoxins of 15 ng/g set by the Codex regulation⁽¹⁷⁾. Considering the tropical weather in Malaysia, products such as peanut stored under this condition are very susceptible to aflatoxins contamination. Regular monitoring of aflatoxins content in peanut is recommended.

ACKNOWLEDGMENTS

The authors would like to acknowledge UPM for sponsoring this research through RUGS fund Project No 02/01/07/0024RU. We are also grateful to Mr. Ismail Muit for his kind help and cooperation in this study.

REFERENCES

- Juan, C., Zinedine, A., Molto, J. C., Idrissi, L. and Manes, J. 2008. Aflatoxins levels in dried fruits and nuts from Rabat-Sale, Morocco. *Food Control* 19: 849-853.
- Bennett, J. W. and Klich, M. 2003. Mycotoxins. *Clin. Microbiol. Rev.* 16: 497-516.
- Akbas, M. Y. and Ozdemir, M. 2006. Effect of different ozone treatments on aflatoxin degradation and physicochemical properties of pistachios. *J. Sci. Food Agric.* 86: 2099-2104.
- Cheraghali, A. M., Yazdanpanah, H., Doraki, N., Abouhossain, G., Hassibi, M., Ali-abadi, S., Aliakbarpoor, M., Amirahmadi, M., Askarian, A., Fallah, N., Hashemi, T., Jalali, M., Kalantari, N., Khodadadi, E., Maddah, B., Mohit, R., Mohseny, M., Phaghihy, Z., Rahmani, A., Setoodeh, L., Soleimany, E. and Zamanian, F. 2007. Incidence of aflatoxins in Iran pistachio nuts. *Food Chem. Toxicol.* 45: 812-816.
- Rustom, I. 1997. Aflatoxin in food and feed: Occurrence, legislation and inactivation by physical methods. *Food Chem.* 59: 57-67.
- Chun, H. S., Kim, H. J., Ok, H. E., Hwang, J. B. and Chung, D. H. 2007. Determination of aflatoxin levels in nuts and their products consumed in South Korea. *Food Chem.* 102: 385-391.
- Yazdanpanah, H., Mohammadi, T., Abouhossain, G. and Cheraghali, A. M. 2005. Effect of roasting on degradation of aflatoxins in contaminated pistachio nuts. *Food Chem. Toxicol.* 43: 1135-1139.
- Dorner, J. W., Cole, R. J., Connick, W. J., Daigle, D. J., McGuire, M. R. and Shasha, B. S. 2003. Evaluation of biological control formulation to reduce aflatoxin contamination in peanuts. *Biol. Control* 26: 318-324.
- Porter, D. M., Wright, F. S. and Steele, J. L. 1986. Relationship to microscopic shell damage to colonization of peanut by *Aspergillus flavus*. *Oleagineux* 41: 23-27.
- Ali, N. 2000. Aflatoxins in Malaysian food. *Mycotoxins* 50: 31-35.
- Council for Agricultural Science and Technology. 2003. Mycotoxins: Risks in Plant, Animal, and Human Systems, Ames, Iowa, U.S.A. <http://www.cast-science.org/>
- Truckness, M. W. 2000. AOAC Official Method 991.31, Aflatoxin in corn, raw peanuts, and peanut butter. *J. AOAC Int.* 49: 22-24.
- Manetta, A. C., Giuseppe, L. D., Giammarco, M., Fusaro, I., Simonella, A., Gramenzi, A. and Formigoni, A. 2005. High performance liquid chromatography with post-column derivatization and fluorescence detection for sensitive determination of aflatoxin M₁ in milk and cheese. *J. Chromatogr. A* 1083: 219-222.
- Papadopoulou-Bouraoui, A., Stroka, J. and Anklam, E. 2000. Comparison of two post-column derivatization systems, ultraviolet irradiation and electrochemical determination, for the liquid chromatographic determination of aflatoxins in food. *J. AOAC Int.* 85: 411-416.
- Moazami, E. F. and Jinap, S. 2009. Natural occurrence of deoxynivalenol (DON) in wheat based noodles consumed in Malaysia. *Microchem. J.* 93: 25-28.
- European Commission. 2006. Commission Regulation No. 401/2006 of 23 February 2006. Laying down the methods of sampling and analysis for the official control of the levels of mycotoxins in foodstuffs. *OJEU*. No. 401. www.icc.or.at/task/EC401-2006.pdf
- CODEX STAN 209-1999, Rev. 1-2001. 2001. Maximum level and sampling plan for total aflatoxins in peanuts intended for further processing. http://siweb.dss.go.th/standard/Fulltext/codex/CXS_209E.pdf
- Ali, N., Hashim, N. H. and Yoshizawa, T. 1999. Evaluation and application of a simple and rapid method for the analysis of aflatoxins in commercial foods from Malaysia and the Philippines. *Food Addit. Contam.* 16: 273-280.
- Haydar, M., Benelli, L. and Brera, C. 1990. Occurrence of aflatoxins in Syrian foods and foodstuffs: A preliminary study. *Food Chem.* 37: 261-268.
- Farombi, E. O. 2006. Contamination of foods in developing countries: Implications for hepatocellular carcinoma and chemopreventive strategies. *Afr. J. Biotechnol.* 5: 1-14.
- Abdulkadar, A. H. W., AL-ALI, A. and Al-Jedah, J. 2000. Aflatoxin contamination in edible nuts imported

- in Qatar. *Food Control* 11: 157-160.
22. Sulaiman, M. R., Yee C. F., Hamid, A. and Yatim, A. M. 2007. The occurrence of aflatoxins in raw shelled peanut samples from three districts of perak, Malaysia. *EJEAFChE* 6: 2045-2052.
 23. Mat Isa, A. and Tee, E. S. 1984. The status of aflatoxin research in Malaysia. In: Country report presented at the First Technical Consultation of ASEAN Mycotoxin Experts. Kuala Lumpur, Malaysia.
 24. Ministry of Health Malaysia. 2006. Food Consumption Statistics of Malaysia. 2002/2003. pp. 7. Food Safety and Quality Division, and Family Health Development Division, Ministry of Health Malaysia, Malaysia.