



2011

## Inhibitory effects of *Polygonum cuspidatum* on the Epstein-Barr virus lytic cycle

Follow this and additional works at: <https://www.jfda-online.com/journal>

---

### Recommended Citation

Yiu, C.-Y.; Chen, S.-Y.; Huang, C.-W.; Yeh, D.-B.; and Lin, T.-P. (2011) "Inhibitory effects of *Polygonum cuspidatum* on the Epstein-Barr virus lytic cycle," *Journal of Food and Drug Analysis*: Vol. 19 : Iss. 2 , Article 3.

Available at: <https://doi.org/10.38212/2224-6614.2234>

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

# Inhibitory Effects of *Polygonum cuspidatum* on the Epstein-Barr Virus Lytic Cycle

CHING-YI YIU<sup>1</sup>, SHIH-YING CHEN<sup>2</sup>, CHING-WEN HUANG<sup>3</sup>, DONG-BOR YEH<sup>3</sup> AND TSUEY-PIN LIN<sup>4\*</sup>

<sup>1</sup> Department of Otolaryngology, Chi Mei Medical Center, Liouying, Tainan, Taiwan R.O.C.

<sup>2</sup> Department of Applied Life Science and Health, Chia-Nan University of Pharmacy and Science, Tainan, Taiwan R.O.C.

<sup>3</sup> Department of Biotechnology, Chia-Nan University of Pharmacy and Science, Tainan, Taiwan R.O.C.

<sup>4</sup> Department of Health and Nutrition, Chia-Nan University of Pharmacy and Science, Tainan, Taiwan R.O.C.

(Received: September 2, 2010; Accepted: December 14, 2010)

## ABSTRACT

*Polygonum cuspidatum* is widely used as a medicinal herb in Asia. In this study, we examined the ethanolic extract of *Polygonum cuspidatum* roots (PcE) for their capacity to inhibit the EBV lytic cycle. The cell viability was determined by the MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] method. The expression of Epstein-Barr virus (EBV) lytic proteins was analyzed by flow cytometric assay. Real-time quantitative PCR was used to measure the EBV virion production. Transient transfection analysis was subsequently performed to assess the transcription of lytic genes, including BRLF1 and BZLF1. Results showed that the PcE inhibited the transcription of EBV immediate early genes, the expression of EBV lytic proteins, including Rta, Zta, and EA-D and reduced the production of virus particles, showing that PcE is useful for preventing the proliferation of the virus.

Key words: *Polygonum cuspidatum*, antiviral activity, EBV

## INTRODUCTION

Epstein-Barr virus (EBV) is a gammaherpesvirus, which infects human lymphoid cells and epithelial cells<sup>(1)</sup>. Infection by this virus is associated with a number of human cancers, including Burkitt's lymphoma<sup>(2)</sup>, nasopharyngeal carcinoma (NPC)<sup>(3)</sup> and Hodgkin's disease<sup>(4)</sup>. The reactivation of EBV from latency to the lytic cycle is necessary for the virus to produce virions to establish infections<sup>(5,6)</sup>. At the onset of the lytic cycle, EBV expresses two transcription factors, Rta and Zta, which are transcribed from BZLF1 and BRLF1, respectively<sup>(7)</sup>. These two proteins trigger an ordered cascade of the expression of viral lytic genes, including that of BMRF1 and BALF5, which encoded diffused early antigen (EA-D) and DNA polymerase<sup>(8)</sup>. Rta and Zta can also activate IL-6 transcription in lytically-infected B cells which leads to immortalized B cells<sup>(9)</sup>. Therefore, an effective strategy to block the viral lytic cycle is of value to reduce the risk of the disease and to improve the clinical outcome. Earlier studies have established that lytic EBV replication is inhibited by acyclovir and ganciclovir, which inhibit specifically the function of viral-encoded DNA polymerase<sup>(10)</sup>. An earlier

study showed that epigallocatechin gallate (EGCG) inhibits the expression of the EBV immediate-early genes transcription at a concentration of 50  $\mu$ M<sup>(11)</sup>. Ethanolic extract from *Andrographis paniculata* and andrographolide also inhibit the expression of Rta, Zta and EA-D at 25  $\mu$ g/mL and 5  $\mu$ g/mL, respectively<sup>(12)</sup>.

*Polygonum cuspidatum* is an herbal medicine, which is commonly used for the treatment of atherosclerosis, as well as other medical ailments including cancers, asthma, hypertension and cough<sup>(13)</sup>. The methanolic extract of *P. cuspidatum* roots contains anthraglycoside B, physcion, piceid, emodin and resveratrol<sup>(14)</sup>. Previous studies showed that the ethanolic extract of *P. cuspidatum* inhibits hepatitis B virus replication<sup>(15)</sup>. In addition, resveratrol, the major compound of *P. cuspidatum*, showed antiviral ability against human cytomegalovirus (HCMV)<sup>(16)</sup>, herpes simplex virus type 1 (HSV-1)<sup>(17)</sup>, influenza virus<sup>(18)</sup>, vaccinia virus<sup>(19)</sup>, EBV<sup>(20)</sup> and Varicella-zoster virus (VZV)<sup>(21)</sup>. Furthermore, resveratrol synergistically enhances the anti-HIV-1 activity of the nucleoside analogues zidovudine (AZT), zalcitabine (ddC) and didanosine (ddI)<sup>(22)</sup>. Emodin has an inhibitory effect on HBV<sup>(23)</sup> and HSV<sup>(24)</sup> replication. Moreover, emodin blocks the SARS coronavirus spike protein and angiotensin-converting enzyme 2 interaction<sup>(25)</sup>. However, no study

\* Author for correspondence. Fax: +886-6-2667327;  
E-mail: tplin007@mail.chna.edu.tw

has been reported on the inhibition of EBV lytic cycle by *P. cuspidatum*. Thus, our study is to evaluate the ethanolic extract of *P. cuspidatum* roots (PcE) on the inhibition of the transcription of EBV immediate early genes, the expression of EBV lytic proteins, including Rta, Zta, and EA-D and virion production.

## MATERIALS AND METHODS

### I. Material

*P. cuspidatum* was collected from the San Dei Men area in Pingtung County, Taiwan and verified by Prof. C. S. Kuoh. The specimen was deposited in the herbarium of the National Cheng Kung University, Tainan, Taiwan.

### II. Extraction of *P. cuspidatum*

Ten grams of dried powder from the *P. cuspidatum* root was extracted three times in 100 mL of ethanol by refluxing at 85°C for 2 h. After each extraction, the ethanol fraction was collected by filtration. The ethanol was then removed by rotary evaporation and a yield of 21.6% was obtained. The resulting residues were finally dissolved in DMSO.

### III. HPLC Analysis

The components in PcE were analyzed by high performance liquid chromatography (HPLC) using a LiChrospher 100 RP-18e (250 × 4.6 mm i.d., 5 μm) column (Merck, Darmstadt, Germany). The mobile phase consisted of (A) water and (B) methanol. Gradient elution was performed as follows: 30 - 50% B in 0 - 15 min, 50 - 90% B in 15 - 35 min, and 95% B in 35 - 45 min. The flow rate was set at 1 mL/min. The effluent was monitored from 210 to 500 nm by a diode array detector.

### IV. Cell Culture and Lytic Induction of EBV

P3HR1, a Burkitt's lymphoma cell line which was latently infected by EBV, was cultured in RPMI 1640 medium containing 10% fetal calf serum (Biological Industries, Israel). Cells were treated with 3 mM of sodium butyrate (SB) to induce the EBV lytic cycle<sup>(26)</sup>.

### V. Cell Viability Assay

A 1 mg/mL solution of [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) in RPMI 1640 medium was added into  $1 \times 10^5$  P3HR1 cells. The dehydrogenase activity of the viable cells was measured using the method of Carmichael *et al*<sup>(27)</sup>.

### VI. Flow Cytometric Analysis

Flow Cytometric Analysis was performed as

described<sup>(20)</sup>, using mouse monoclonal anti-Rta, anti-Zta (Argene, Varilhes, France), and anti-EA-D antibodies (Chemicon, Temecula, CA, USA). Secondary antibodies used in the study included Alexa Fluor 488-conjugated goat anti-mouse IgG from Invitrogen. Finally, the cells were resuspended in 1% paraformaldehyde and analyzed using a flow cytometer (Model FACScanTO, BD Biosciences)

### VII. Transient Transfection and Luciferase Assays

P3HR1 cells ( $5 \times 10^6$ ) were transfected with 10 μg of pRLUC and pZLUC with a Bio-Rad electroporator using the method of Chang *et al*<sup>(28,29)</sup>. The cells were harvested at 24 h after transfection, and luciferase activity was determined using a luminometer (Berthold, Germany)<sup>(28)</sup>.

### VIII. Quantification of EBV Particles Produced by P3HR1 Cells

P3HR1 cells were cultured for 5 days. EBV particles released into the culture medium were harvested by centrifugation. Viral DNA was extracted and the amount of EBV DNA was determined by real-time PCR using an iCyclerIQ multicolor real-time PCR detection system (BioRad, CA, USA) with primer and a probe that were specific to the EBNA1 gene<sup>(30)</sup>. The copy number of the EBV genome was calculated using maxi-EBV DNA extracted from *E. coli* as a reference. The molecular weight of maxi-EBV is about  $1.2 \times 10^7$  and 1 ng of maxi-EBV equals to  $5.05 \times 10^6$  copies of the maxi-EBV genome.

### IX. Statistical Analysis

Data were analyzed statistically by one-way analysis of variance (ANOVA) using the SAS JMP 6.0 software package. Data are presented as means ± SD and a *p* value of < 0.05 was regarded as significant.


## RESULTS

### I. Identification of Resveratrol and Emodin in PcE

The PcE was analyzed by HPLC and the identification of the target compounds in PcE (resveratrol and emodin) was based on retention time and UV spectra, compared against those of the pure standards of resveratrol and emodin. The peak eluting at 10.99 min and 33.01 min were ascribed to resveratrol and emodin, respectively (Figure 1).

### II. Determining the Toxicity of PcE to P3HR1 Cells

PcE was added into  $1 \times 10^5$  cells/mL. After 24 h of treatment, cell viability was determined by MTT assay. The results showed that PcE affected the viability of P3HR1 cells in a dose-dependent manner. PcE decreased cell viability to 50% (CC<sub>50</sub>) at 84.56 μg/mL (Figure 2).



**Figure 1.** Chromatogram of the ethanolic extract of *Polygonum cuspidatum* root (PcE) by HPLC. Mobile phase: methanol-water (methanol: 0-15 min, 30-50%; 15-35 min, 50-90%; 35-45 min, 95%); flow rate: 1.0 mL/min; detection wavelength: 280 nm.

### III. Flow Cytometric Analysis of the Expression of EBV Lytic Proteins


The presence of EBV lytic proteins in P3HR1 cells was further analyzed by flow cytometric analysis. The P3HR1 population that was untreated with SB expressed Rta, Zta and EA-D was 2.9%, 2.9% and 4.8%, respectively (Figure 3A). The population that expressed three proteins after SB treatment increased to 49.2%, 51.4% and 43%, respectively. When 12.5  $\mu\text{g/mL}$  of PcE was added before lytic induction, the population that expressed Rta, Zta and EA-D decreased to 34.5%, 29.4% and 38.6%, respectively (Figure 3A). The population that expressed Rta, Zta, and EA-D further decreased when the concentration of PcE increased to 25 and 50  $\mu\text{g/mL}$ . At 50  $\mu\text{g/mL}$ , the population that expressed Rta, Zta, and EA-D decreased to 7.8%, 7.3%, and 1%, respectively (Figure 3A). These results showed that PcE inhibited the expression of Rta, Zta and EA-D in a dose-dependent manner (Figure 3B). The concentration of PcE required to inhibit EBV immediate-early proteins expression by 50% ( $\text{EC}_{50}$ ) is approximately 29  $\mu\text{g/mL}$ .

### IV. Inhibiting the Transcription of EBV Immediate-Early Genes

The activity of the BRLF1 and BZLF1 promoters was analyzed by transient transfection assay in P3HR1 cells using reporter plasmids, pRLUC and pZLUC, respectively. The results showed that 12.5, 25 and 50  $\mu\text{g/mL}$  of PcE inhibited the BRLF1 promoter activity (35%, 54% and 77%, respectively). At these concentrations, PcE also inhibited BZLF1 promoter activity (45%, 63% and 70%, respectively) (Figure 4).

### V. Inhibition of the Production of EBV Particles

P3HR1 cells were treated with 12.5 - 50  $\mu\text{g/mL}$  of PcE




**Figure 2.** Effect of PcE on cell viability. P3HR1 cells ( $1 \times 10^5$  cell/mL) were treated with PcE. Cell numbers and viability were monitored by MTT assay after 24 h of treatment. The data are presented as mean  $\pm$  SD from three independent experiments. At 50 and 100  $\mu\text{g/mL}$ ,  $p < 0.05$  vs. control.

before lytic induction. After culturing for five days, EBV particles released into the medium were isolated. Real-time qPCR was used to determine the amount of EBV DNA purified from the viral particles. The results showed that PcE at a concentration of 25  $\mu\text{g/mL}$  decreased virus production by 76% (Figure 5). The effective concentration of PcE that inhibited EBV genome copy numbers by 50% ( $\text{EC}_{50}$ ) is 17.9  $\mu\text{g/mL}$ .

## DISCUSSION

Previous studies showed that lytic EBV proteins actually induce the expression of B-cell growth factor, IL-6, cellular IL-10 and viral IL-10, allowing B cells to grow




**Figure 3.** Inhibition of the expression of EBV lytic proteins byPcE. P3HR1 cells were untreated (latent) or treated with PcE (0, 12.5, 25 and 50 µg/mL) before SB induction (A). At 24 h after treatment, cells were stained with primary mouse monoclonal antibodies (anti-Rta or anti-Zta or anti-EA-D) and Alexa Fluor 488-conjugated goat anti-mouse IgG and then analyzed by flow cytometry. M2 represents fluorescence stained cells. The number of cells expressing EBV lytic proteins was quantified by averaging the results from three independent experiments (B). The data are presented as means with standard deviation. EBV Lytic protein expression (%) = (M2 of w/wo PcE treated before SB induction - M2 of Latent/M2 of SB treated - M2 of Latent) × 100%.


more efficiently<sup>(31,32)</sup>. Lytically-infected cells also produce VEGF and thus contribute to angiogenesis in both B-cell and epithelial-cell malignancies<sup>(32)</sup>. Therefore, new treatment strategies aimed at completely suppressing the expression of all lytic viral proteins are useful in controlling early EBV-associated malignancies.

In this study, it was found that PcE significantly reduces the expression of EBV immediate-early proteins, Rta, Zta

and EA-D in a dose-dependent manner (Figure 3). In other words, PcE interferes with an early step of the EBV replication cycle. The concentration of PcE required to inhibit EBV immediate-early proteins expression by 50% (EC<sub>50</sub>) is approximately 29 µg/mL. Moreover, PcE exhibits cytotoxicity in P3HR1 cells and the CC<sub>50</sub> is 84.56 µg/mL. Our study also demonstrates that PcE inhibits the transcriptional activity of BRLF1 and BZLF1 promoters, which in turn,



**Figure 4.** Inhibition of the transcription of the BRLF1 and BZLF1 promoters by PcE. P3HR1 cells were transfected with pRLUC or pZLUC and treated with PcE before lytic induction. At 24 h after transfection, luciferase activity was measured using a luminometer. Each transfection experiment was performed three times, and each sample in the experiment was prepared in duplicate.



**Figure 5.** Effect of PcE on EBV particles production. P3HR1 cells were treated with PcE in the presence of sodium butyrate (SB). Five days after the treatment, EBV particles were harvested by centrifugation. The amounts of EBV DNA isolated from the particles were determined by real-time qPCR. The copy number of EBV genome was calculated by using maxi-EBV. EBV Genome (%) =  $\frac{\text{Log of copy number (PcE treated before SB induction)} - \text{PcE and SB untreated}}{\text{Log of copy number (SB treated only)} - \text{PcE and SB untreated}} \times 100\%$ . The data are presented as mean  $\pm$  SD from at least three independent experiments.

affects viral lytic proteins expression. In addition, the inhibition actually decreases the production of mature viral particle. Real-time qPCR indicates that the effective concentration of PcE that inhibits EBV genome copy numbers by 50% ( $EC_{50}$ ) is 17.9  $\mu\text{g/mL}$  (Figure 5), compared to that obtained by flow cytometry (29  $\mu\text{g/mL}$ ). Besides Rta and Zta proteins, the activation of EBV lytic cycle requires other factors, such as the

MBD1-containing chromatin-associated factor 1 (MCAF1)<sup>(33)</sup>. These data implied that PcE may inhibit other lytic genes involved in EBV lytic cycle. However, the inhibitory profile remains unchanged, indicating that flow cytometry is as reliable as the real-time qPCR method for initial drug screening.

The molecular mechanism underlying the inhibition of EBV early gene expression by PcE is unclear. Results showed that PcE contains resveratrol and emodin (Figure 1). Previous studies showed that emodin and resveratrol inhibit the activation of p38 MAPK, ERK and JNK signaling and affect the activation of the promoters that are activated by AP-1 or ATF2<sup>(34)</sup>, as both the BRLF1 and BZLF1 promoters are strongly activated by AP-1 and ATF2<sup>(35-37)</sup>. The transcriptional inhibition of BRLF1 and BZLF1 promoters may be due to resveratrol that activates the deacetylase activity of sirtuin protein<sup>(20)</sup>. Therefore, we suggest that emodin and resveratrol may be involved in the inhibition of PcE in the EBV lytic cycle.

## CONCLUSIONS

In this study, the results clearly demonstrated that PcE inhibits the transcription of lytic genes and the lytic cycle of EBV to reduce the production of viral particles. PcE could potentially be used for the development of anti-EBV drugs.

## ACKNOWLEDGMENTS

This work was supported by research grants 96-2320-B-041-009 from the National Science Council, Taiwan, and CLFHR9620 from Chi Mei Medical Center, Liouying, Taiwan.

## REFERENCES

- Roizman, B., Carmichael, L. E., Deinhardt, F., de-The, G., Nahmias, A. J., Plowright, W., Rapp, F., Sheldrick, P., Takahashi, M. and Wolf, K. 1981. Herpesviridae. Definition, provisional nomenclature, and taxonomy. The Herpesvirus Study Group, the International Committee on Taxonomy of Viruses. *Intervirology*. 16: 201-217.
- Magrath, I., Jain, V. and Bhatia, K. 1992. Epstein-Barr virus and Burkitt's lymphoma. *Semin. Cancer Biol.* 3: 285-295.
- zur Hausen, H., Schulte-Holthausen, H., Klein, G., Henle, W., Henle, G., Clifford, P. and Santesson, L. 1970. EBV DNA in biopsies of Burkitt tumours and anaplastic carcinomas of the nasopharynx. *Nature*. 228: 1056-1058.
- Weiss, L. M., Movahed, L. A., Warnke, R. A. and Sklar, J. 1989. Detection of Epstein-Barr viral genomes in Reed-Sternberg cells of Hodgkin's disease. *N. Engl. J. Med.* 320: 502-506.
- Schwarzmann, F., Jager, M., Prang, N. and Wolf, H. 1998. The control of lytic replication of Epstein-Barr virus in B lymphocytes (Review). *Int. J. Mol. Med.* 1: 137-142.
- Hopwood, P. A., Brooks, L., Parratt, R., Hunt, B. J., Bokhari, M., Thomas, J. A., Yacoub, M. and Crawford, D. H. 2002. Persistent Epstein-Barr virus infection: unrestricted latent and lytic viral gene expression in healthy immunosuppressed transplant recipients. *Transplantation*. 74: 194-202.
- Giot, J. F., Mikaelian, I., Buisson, M., Manet, E., Joab, I., Nicolas, J. C. and Sergeant, A. 1991. Transcriptional interference between the EBV transcription factors EB1 and R: both DNA-binding and activation domains of EB1 are required. *Nucleic Acids Res.* 19: 1251-1258.
- Fixman, E. D., Hayward, G. S. and Hayward, S. D. 1992. trans-acting requirements for replication of Epstein-Barr virus ori-Lyt. *J. Virol.* 66: 5030-5039.
- Tosato, G., Tanner, J., Jones, K. D., Revel, M. and Pike, S. E. 1990. Identification of interleukin-6 as an autocrine growth factor for Epstein-Barr virus-immortalized B cells. *J. Virol.* 64: 3033-3041.
- Meerbach, A., Holy, A., Wutzler, P., De Clercq, E. and Neyts, J. 1998. Inhibitory effects of novel nucleoside and nucleotide analogues on Epstein-Barr virus replication. *Antivir. Chem. Chemother.* 9: 275-282.
- Chang, L. K., Wei, T. T., Chiu, Y. F., Tung, C. P., Chuang, J. Y., Hung, S. K., Li, C. and Liu, S. T. 2003. Inhibition of Epstein-Barr virus lytic cycle by (-)-epigallocatechin gallate. *Biochem. Biophys. Res. Commun.* 301: 1062-1068.
- Lin, T. P., Chen, S. Y., Duh, P. D., Chang, L. K. and Liu, Y. N. 2008. Inhibition of the Epstein-Barr virus lytic cycle by andrographolide. *Biol. Pharm. Bull.* 31: 2018-2023.
- Yi, T., Zhang, H. and Cai, Z. 2007. Analysis of *Rhizoma Polygoni Cuspidati* by HPLC and HPLC-ESI/MS. *Phytochem. Anal.* 18: 387-392.
- Chu, X., Sun, A. and Liu, R. 2005. Preparative isolation and purification of five compounds from the Chinese medicinal herb *Polygonum cuspidatum* Sieb. et Zucc by high-speed counter-current chromatography. *J. Chromatogr. A*. 1097: 33-39.
- Chang, J. S., Liu, H. W., Wang, K. C., Chen, M. C., Chiang, L. C., Hua, Y. C. and Lin, C. C. 2005. Ethanol extract of *Polygonum cuspidatum* inhibits hepatitis B virus in a stable HBV-producing cell line. *Antiviral Res.* 66: 29-34.
- Evers, D. L., Wang, X., Huong, S. M., Huang, D. Y. and Huang, E. S. 2004. 3,4',5-Trihydroxy-trans-stilbene (resveratrol) inhibits human cytomegalovirus replication and virus-induced cellular signaling. *Antiviral Res.* 63: 85-95.
- Docherty, J. J., Fu, M. M., Stiffler, B. S., Limperos, R. J., Pokabla, C. M. and DeLucia, A. L. 1999. Resveratrol inhibition of herpes simplex virus replication. *Antiviral Res.* 43: 145-155.
- Kim, Y., Narayanan, S. and Chang, K. O. 2010. Inhibition of influenza virus replication by plant-derived isoquercetin. *Antiviral Res.* 88: 227-235.
- Cheltsov, A. V., Aoyagi, M., Aleshin, A., Yu, E. C., Gilliland, T., Zhai, D., Bobkov, A. A., Reed, J. C., Liddington, R. C. and Abagyan, R. 2010. Vaccinia virus virulence factor NIL is a novel promising target for antiviral therapeutic intervention. *J. Med. Chem.* 53: 3899-3906.
- Yiu, C. Y., Chen, S. Y., Chang, L. K., Chiu, Y. F. and Lin, T. P. 2010. Inhibitory effects of resveratrol on the Epstein-Barr virus lytic cycle. *Molecules*. 15: 7115-7124.
- Docherty, J. J., Sweet, T. J., Bailey, E., Faith, S. A. and Booth, T. 2006. Resveratrol inhibition of varicella-zoster virus replication in vitro. *Antiviral Res.* 72: 171-177.
- Heredia, A., Davis, C. and Redfield, R. 2000. Synergistic inhibition of HIV-1 in activated and resting peripheral blood mononuclear cells, monocyte-derived macrophages, and selected drug-resistant isolates with nucleoside analogues combined with a natural product, resveratrol. *J. Acquir. Immune Defic. Syndr.* 25: 246-255.
- Shuangsoo, D., Zhengguo, Z., Yunru, C., Xin, Z., Baofeng, W., Lichao, Y. and Yan'an, C. 2006. Inhibition of the replication of hepatitis B virus in vitro by emodin. *Med. Sci. Monit.* 12: BR302-306.
- Hsiang, C. Y. and Ho, T. Y. 2008. Emodin is a novel alkaline nuclease inhibitor that suppresses herpes simplex virus type 1 yields in cell cultures. *Br. J. Pharmacol.* 155: 227-235.
- Ho, T. Y., Wu, S. L., Chen, J. C., Li, C. C. and Hsiang, C. Y. 2007. Emodin blocks the SARS coronavirus spike protein and angiotensin-converting enzyme 2 interaction. *Antiviral Res.* 74: 92-101.
- Luka, J., Kallin, B. and Klein, G. 1979. Induction of the Epstein-Barr virus (EBV) cycle in latently infected cells by n-butyrate. *Virology*. 94: 228-231.
- Carmichael, J., DeGraff, W. G., Gazdar, A. F., Minna, J. D. and Mitchell, J. B. 1987. Evaluation of a tetrazolium-based semiautomated colorimetric assay: assessment of chemosensitivity testing. *Cancer Res.* 47: 936-942.

28. Chang, L. K. and Liu, S. T. 2000. Activation of the BRLF1 promoter and lytic cycle of Epstein-Barr virus by histone acetylation. *Nucleic Acids Res.* 28: 3918-3925.
29. Chang, P. J., Chang, Y. S. and Liu, S. T. 1998. Role of Rta in the translation of bicistronic BZLF1 of Epstein-Barr virus. *J. Virol.* 72: 5128-5136.
30. Chiu, Y. F., Tung, C. P., Lee, Y. H., Wang, W. H., Li, C., Hung, J. Y., Wang, C. Y., Kawaguchi, Y. and Liu, S. T. 2007. A comprehensive library of mutations of Epstein Barr virus. *J. Gen. Virol.* 88: 2463-2472.
31. Jones, R. J., Seaman, W. T., Feng, W. H., Barlow, E., Dickerson, S., Delecluse, H. J. and Kenney, S. C. 2007. Roles of lytic viral infection and IL-6 in early versus late passage lymphoblastoid cell lines and EBV-associated lymphoproliferative disease. *Int. J. Cancer.* 121: 1274-1281.
32. Hong, G. K., Kumar, P., Wang, L., Damania, B., Gulley, M. L., Delecluse, H. J., Polverini, P. J. and Kenney, S. C. 2005. Epstein-Barr virus lytic infection is required for efficient production of the angiogenesis factor vascular endothelial growth factor in lymphoblastoid cell lines. *J. Virol.* 79: 13984-13992.
33. Chang, L. K., Chung, J. Y., Hong, Y. R., Ichimura, T., Nakao, M. and Liu, S. T. 2005. Activation of Sp1-mediated transcription by Rta of Epstein-Barr virus via an interaction with MCAF1. *Nucleic Acids Res.* 33: 6528-6539.
34. Kang, O. H., Jang, H. J., Chae, H. S., Oh, Y. C., Choi, J. G., Lee, Y. S., Kim, J. H., Kim, Y. C., Sohn, D. H., Park, H. and Kwon, D. Y. 2009. Anti-inflammatory mechanisms of resveratrol in activated HMC-1 cells: pivotal roles of NF-kappaB and MAPK. *Pharmacol. Res.* 59: 330-337.
35. Matusali, G., Arena, G., De Leo, A., Di Renzo, L. and Mattia, E. 2009. Inhibition of p38 MAP kinase pathway induces apoptosis and prevents Epstein Barr virus reactivation in Raji cells exposed to lytic cycle inducing compounds. *Mol. Cancer.* 8: 18.
36. Rahman, M. M., Kukita, A., Kukita, T., Shobuike, T., Nakamura, T. and Kohashi, O. 2003. Two histone deacetylase inhibitors, trichostatin A and sodium butyrate, suppress differentiation into osteoclasts but not into macrophages. *Blood.* 101: 3451-3459.
37. Adamson, A. L., Darr, D., Holley-Guthrie, E., Johnson, R. A., Mauser, A., Swenson, J. and Kenney, S. 2000. Epstein-Barr virus immediate-early proteins BZLF1 and BRLF1 activate the ATF2 transcription factor by increasing the levels of phosphorylated p38 and c-Jun N-terminal kinases. *J. Virol.* 74: 1224-1233.