



2009

Sequence analysis based on ITS1 region of nuclear ribosomal DNA of *Amomum villosum* and ten species of *Alpinia*

Follow this and additional works at: <https://www.jfda-online.com/journal>

Recommended Citation

Qiao, C.-F.; Han, Q.-B.; Zhao, Z.-L.; Wang, Z.-T.; Xu, L.-S.; and Xu, H.-X. (2009) "Sequence analysis based on ITS1 region of nuclear ribosomal DNA of *Amomum villosum* and ten species of *Alpinia*," *Journal of Food and Drug Analysis*: Vol. 17 : Iss. 2 , Article 1.

Available at: <https://doi.org/10.38212/2224-6614.2288>

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

Sequence Analysis Based on ITS1 Region of Nuclear Ribosomal DNA of *Amomum villosum* and Ten Species of *Alpinia*

CHUN-FENG QIAO¹, QUAN-BIN HAN¹, ZHI-LI ZHAO³, ZHENG-TAO WANG²,
LUO-SHAN XU^{2*} AND HONG-XI XU^{1*}

¹ Chinese Medicine Laboratory, Hong Kong Jockey Club Institute of Chinese Medicine, Hong Kong, China

² Department of Pharmacognosy, China Pharmaceutical University, Nanjing, China

³ Department of Chinese Material Medica, Shanghai University of Traditional Chinese Medicine, Shanghai, China

(Received: November 19, 2008; Accepted: February 3, 2009)

ABSTRACT

Ripe fruits of *Amomum villosum* Lour. are important traditional Chinese medicines and food spices. Due to similar morphological and anatomical characteristics, fruits from some *Alpinia* species are used clinically as substitutes or adulterants of Fructus Amomi. In this study, we determined the nrDNA ITS1 (internal transcribed space 1) sequences from *Amomum villosum* and ten *Alpinia* species, which are adulterants of Fructus Amomi. The ITS1 sequences of these species were 175~179 bp in length, and the G+C contents were 53.37~56.74%. Sequence alignment revealed that there were 55 polymorphic nucleotide sites among the tested species. The base substitution were 0.57~9.60% within the ten *Alpinia* species, but 17.61~21.59% between *Amomum villosum* and the *Alpinia* species. These molecular data indicated that ITS1 sequences can be used to differentiate Fructus Amomi and its substitutes or adulterants from various *Alpinia* species.

Key words: *Amomum villosum*, Fructus Amomi, *Alpinia*, nrDNA ITS1

INTRODUCTION

Amomum villosum Lour. (Zingiberaceae) is an aromatic plant distributed mainly in Guangdong, Yunnan, Fujian and Guangxi Province of China⁽¹⁾. The ripe fruits of *A. villosum*, named "Sha-ren" in Chinese, have been used as traditional medicines and food spices in China for centuries⁽²⁾. Its main function is to treat dyspepsia, gastric disease, vomiting and diarrhea⁽³⁾. Experimental bioassays have demonstrated that the extracts of Fructus Amomi help improving the gastrointestinal milieu⁽⁴⁾ and show obvious analgesic and anti-inflammatory effects^(5,6).

According to the Chinese Pharmacopoeia (2005), Fructus Amomi is derived from *A. villosum*⁽³⁾. However, our literature search and survey of the commercial crude drugs of Fructus Amomi revealed that the fruits of some *Alpinia* species from the same family, such as *A. japonica*, *A. pinnanensis* and *A. guinanensis*, are also used clinically as substitutes or adulterants in China⁽⁷⁻⁹⁾. However, improper usage may cause inconsistent thera-

peutic effects, or even jeopardize the safety of consumers. Therefore, accurate identification of Fructus Amomi is important for its safety and efficacy.

The morphological and anatomical characteristics of the fruits from these *Alpinia* and Fructus Amomi species are very similar^(10,11). Since the chemical constituents of these species are complicated and still unknown, a chemical analysis method for the quality control of Fructus Amomi has not yet been developed. In recent years, the DNA-based polymorphism assay has been more closely adapted for the identification of herbal medicines, and developed to be a more effective, accurate, reliable and sensitive technology^(12,13). Samples of *A. villosum* from different habitats and adulterants have been analyzed by RAPD (random amplified polymorphic DNA)^(14,15). Furthermore, the ITS (internal transcribed space) sequence of *A. villosum* was also amplified and determined. The results showed that the ITS1 sequence can be used to confirm its production area and determine its adulterants from *Amomum villosum* var. *xanthioides* and *Amomum longiligulare*^(16,17). In the present study, we determined and compared the ITS1 sequences amplified from *A. villosum* and ten *Alpinia* species, which are adulterants of Fructus Amomi, the results indicated that

* Author for correspondence.

LUO-SHAN XU-Tel: +86-25-85391246; Fax: +86-25-85309639;

HONG-XI XU-Tel: +852-34062875; Fax: +852-35517333;

E-mail: xuhongxi@hkjcicm.org

several polymorphic sites may be used as DNA markers for the identification of this herbal medicine.

MATERIALS AND METHODS

I. Materials

The plants of *A. villosum* and ten *Alpinia* species were collected from different regions in China (Table 1). The species were identified carefully according to the morphological characteristics. Voucher samples were deposited in Department of Pharmacognosy, China Pharmaceutical University.

II. DNA Extraction and Purification

The air-dried leaves of the samples were grounded into powder with mortar and pestle in liquid nitrogen. Total DNA was extracted from 100 mg of powdered sample using a modified CTAB (cetyltrimethylammonium bromide) procedure⁽¹⁸⁾ and 1.2 mL of CTAB extraction buffer (1% CTAB; 50 mM Tris-HCl, pH 8.0; 10 mM EDTA; 0.7 M NaCl; 0.14% 2-mercaptoethanol) was added. The suspension was incubated in a water bath at 56°C for 30 min with occasional shaking. After centrifugation, the supernatant was extracted with equal volume of chloroform/isoamyl alcohol (24:1). After centrifugation at 12000 rpm for 15 min, double volume of absolute alcohol was added to the upper aqueous layer. After centrifugation at 12000 rpm for 15 min, the liquid was discarded. Then, the deposit was washed with 70% ethanol twice, air-dried and dissolved with 55 µL double-distilled water for the next process of purification. The total DNA preparation was purified using a PCR Preps DNA Purification

Kit (Watson, China) according to the protocol supplied by the manufacturer.

III. PCR Amplification

The primers used for PCR amplification were designed based on the primers for *Diseae* species (Orchidaceae)⁽¹⁹⁾ and the known rDNA ITS sequences of Zingiberaceae plants. Nucleotide sequences of the primers were as follows: 5'-ATTGAATGGTCCGGTGAAGT-GTTCG-3' (P1) and 5'-AATCCCCGGTTCGCTCGCC-GTTAC-3' (P2). The forward primer P1 was located at the flanking region of the 3' end of 18S and the reverse primer P2 was located at the flanking region of the 5' end of 26S. PCR amplification was performed using 70-80 ng of total DNA as a template in 30 µL of reaction mixture, consisting of 3.0 µL 10 × PCR buffer, 0.15 mM of dNTPs (Sangon, China), 0.20 mM of each primer, and 1.0 U of Taq Polymerase (Promega, USA). Amplification was carried out under the following conditions: pre-cycling at 94°C for 4 min, 30 cycles of denaturation at 94°C for 40 s, annealing at 55°C for 1 min and elongation at 72°C for 3 min, and then final elongation at 72°C for 3 min. PCR products were detected by 1.5% agarose gel (containing 1 µg/mL ethidium bromide) electrophoresis.

IV. Sequencing

The purified PCR products by a PCR Preps DNA Purification Kit (Watson, China) served as the template. The sequencing reaction was performed using the Big DyeTM Terminator RR Mix (Perkin Elmer, USA) with the same primers as in PCR amplification. The thermal cycling conditions was 95°C for 30 s, 35 cycles of 96°C for 10 s, 50°C for 5 s and 60°C for 4 min. After puri-

Table 1. Samples used in DNA sequence analysis

Code	Species	Voucher	Locality	GenBank accession
1	<i>Amomum villosum</i> Lour.	ZHAO97011	Yangchun, Guangdong	EF488008
2	<i>Alpinia japonica</i> (Thunb.) Miq.	ZHAO98004	Conghua, Guangdong	AF254474
3	<i>Alpinia pinnanensis</i> T. L. Wu & Senjen	ZHAO98005	Wuming, Guangxi	AF254470
4	<i>Alpinia guinanensis</i> D. Fang & X. X. Chen	ZHAO98021	Nanning, Guangxi	AF254477
5	<i>Alpinia tonkinensis</i> Gagnep.	ZHAO98011	Nanning, Guangxi	AF254473
6	<i>Alpinia blepharocalyx</i> K. Schum. var. <i>glabrior</i> (Hand-Mazz.) T. L. Wu	ZHAO98006	Wuming, Guangxi	
7	<i>Alpinia suishaensis</i> Hayata	ZHAO98002	Conghua, Guangdong	AF254465
8	<i>Alpinia stachyoides</i> Hance	ZHAO98015	Yangchun, Guangdong	
9	<i>Alpinia polyantha</i> D. Fang	ZHAO98023	Guangzhou, Guangdong	AF254475
10	<i>Alpinia pumila</i> Hook. f.	ZHAO98017	Guangzhou, Guangdong	AF254472
11	<i>Alpinia jianganfeng</i> T. L. Wu	ZHAO97002	Guangzhou, Guangdong	EF488011

fication, the sequencing products were analyzed on an automated DNA sequencer (ABI PRISM™ 310 Genetic Analyzer, USA). The obtained sequences were edited and aligned with the Clustal W program. The nucleotides sequence data of ITS1-5.8S-ITS2 region were deposited in the GenBank nucleotide sequence database with the accession numbers shown in Table 1. The borders of ITS1, 5.8S, ITS2 regions were determined by comparison with the known sequences of Zingiberaceae plants.

RESULTS AND DISCUSSION

A comparison of alignments of the ITS1 sequences from *Amomum villosum* and ten *Alpinia* species is shown

in Figure 1. The lengths and base compositions of the sequences were measured, and the numbers of transitions/transversions and substitution percentage were summarized.

For all the species, the lengths of the ITS1 sequences were 175~179 bp. The G+C contents of the sequences from ten *Alpinia* species were 53.37~55.93%, and that of *Amomum villosum* was slightly higher at 56.74%. The *Alpinia* species are more closely related to each other than to *Amomum villosum*. In fact, the minimum difference (base substitution percentage) between these ten *Alpinia* species was 0.57% (between *A. japonica* and *A. pumila*), and even the maximum was only 9.60% (between *A. tonkinensis* and *A. jianganfeng*). On the other hand, the difference between *Amomum villosum* and the *Alpinia*

	1						70
1	TTGTGAGAGC	ACTGAACAAT	GGATGATTGT	GAACGTGTCA	ACATGCCCT	CTACTTGGCC	CCATTTT--G
2	...A.....	.T....G.CG....	...T.....	..G.....	T.TG...-..G.....
3	...A.....	.T....G.CGA...	...T.....	..G.....	T.TG...-..	..T..G.....
4	...A.....	.T....G.CGC...	...T.....	..GC.....	T.CGC...-..G.-...
5	...A.....	.T..G..G.CGC...	...T.....	..GC.....	A.CGC...-..G.....
6	..A.....	.T....G..GC...	...T.....	..GC.....	T.CGC...-..G..GT.
7	...A.....	.T....G.C	.A...C...	...T.....	..G.....	T.CGC...-..G.....
8	...A.....	.T....G.C	.A...G....	...T.....	..G.....	T.TG...-..G.....
9	...A.....	.T....G.CGC...	...T.....	..G.....	T.TG...-..G.....
10	...A.....	.T....G.C-....	...T.....	..G.....	T.TG...-..G.....
11	...A.....	.T....G.CT.....	..G...A..	T.TG...-..G.....
	71						140
1	GTGGCCGACT	GACCATAGCT	CGGTGCGATC	GGCACCAAGG	AACAATGAAC	TCAGAAGGAG	AGGG-CCCTC
2	.CA....T.G....	T.....-
3	.CA....T.G....	T.....-
4	.CA.TT..T.	..T.G....	T.....T	..T.....
5	.CA.TT..T.	..T.G....	T.....T	..G.....
6	.CA.TT..T.	..T.G....	T.....T	..T...T
7	.CA..T..T.G....	T.....
8	.CA..T..T.	..T.G....	T.....
9	.CA..T..T.G....	T.....
10	.CA....T.G....	T.....A.....
11	.CA....T.G....	T.....
	141			183			
1	GGTGTGCCCG	GGG-AGCCCA	CT-GCGTCAG	AGACGGTCGG	AAT		
2	A.C...G..	A....G...	A...A..G.	...T.CCTCA	...		
3	A.C...G..	A....G...	A...A..G.	...T.CCTCA	...		
4	A.C...G..G...	G...A..G.	...T..CTCA	...		
5	A.C...TG..G...	A...A..G.	...T..CTCA	...		
6	A.C...G.A	...-...G...	A...A..G.	...T.CCTCA	...		
7	A.C...G..	A....G...	A...A..G.	...T.CCTCA	...		
8	A.C...G..G...	A.T..A..G.	...T.CCTCA	...		
9	..C...GT.	A..G..G...	A...A..G.	...T.CCTCA	...		
10	A.C...G..	A....G...	A...A..G.	...T.CCTCA	...		
11	A.C...G..	A....G...	A...A..G.	C..T.CCTCA	C..		

Figure 1. Alignments of ITS1 sequences of *Amomum villosum* and ten *Alpinia* species. Dots represent sequence identity with sample 1, and dashes indicate deletions.

1. *Amomum villosum*; 2. *Alpinia japonica*; 3. *Alpinia pinnanensis*; 4. *Alpinia guinanensis*; 5. *Alpinia tonkinensis*; 6. *Alpinia blepharocalyx* var. *glabrior*; 7. *Alpinia suishaensis*; 8. *Alpinia stachyoides*; 9. *Alpinia polyantha*; 10. *Alpinia pumila*; 11. *Alpinia jianganfeng*

species was 17.61~21.59%, which was obviously much greater. As demonstrated in these molecular data, the ITS1 sequences may serve as a useful tool for comparison between intra-genus or inter-genus samples.

Sequence alignment of the ITS1 from these eleven species revealed that there are 55 polymorphic nucleotide sites and almost every species had their own specific variable bases. Therefore, these polymorphic sites can be used as DNA markers not only to distinguish *Amomum villosum* from *Alpinia* species but also to identify the original species of the raw material.

For correct identification of biological materials, DNA profiling has several advantages over morphological and chemical analyses because genotypes rather than phenotypes are directly assayed; consequently, the result is not affected by environmental factors. The present investigation indicates that DNA authentication is reliable in differentiating *Amomum villosum* from the ten *Alpinia* species. The selected ITS1 sequences may be useful to identify Fructus Amomi and its substitutes or adulterants from various *Alpinia* species. Based on our findings, the prospects of establishing methods to identify the crude drugs of Fructus Amomi are promising.

ACKNOWLEDGEMENTS

We thank Professor Xiao-Bo Li, Shanghai Jiao Tong University and Professor Kai-Ya Zhou, Prof. Yi-Quan Wang and Professor Zhong-Quan Liu, Nanjing Normal University for their assistance in conducting the experiments.

REFERENCES

1. Wu, D. L. 1981. Zingiberaceae. In "Flora Reipublicae Popularis Sinicae". 1st ed. Vol. 16 (2). p. 125. Edition committee of Flora Reipublicae Popularis Sinicae ed. Science Press. Beijing, China.
2. Mei, Q. C. 1994. Fructus Amomum. In "Species Systematization and Quality Evaluation of Commonly Used Chinese Traditional Drugs". 1st ed. Vol. 1. pp. 606-647. Xu, G. J. and Xu, L. S. ed. Fujian Science and Technology Press. Fuzhou, China.
3. Chinese Pharmacopoeia Commission. 2005. Pharmacopoeia of the People's Republic of China (English version), Vol. 1. p. 80. People's Medical Publishing House. Beijing, China.
4. Huang, Y. L., Yen, G. C., Sheu, F., Lin, J. Y. and Chau, C. F. 2007. Dose effects of the food spice cardamom on aspects of hamster gut physiology. Mol. Nutr. Food Res. 51: 602-608.
5. Wu, X. S., Li, X. G., Xiao, F., Zhang, Z. D., Xu, Z. X. and Wang, H. 2004. Studies on the analgesic and anti-inflammatory effect of bornyl acetate in volatile oil from *Amomum villosum*. J. Chin. Med. Mater. 27: 438-439.
6. Wu, X. S., Xiao, F., Zhang, Z. D., Li, X. G. and Xu, Z. X. 2005. Research on the analgesic effect and mechanism of bornyl acetate in volatile oil from *Amomum villosum*. J. Chin. Med. Mater. 28: 505-507.
7. Jia, M. R., Wei, Y. F., Ma, Y. Y., Zhong, G. Y., Zhang, Y. Kuang, T. L. and Huang, X. H. 1985. Research on medicinal "Tu-Sha-Ren" used in Leshan district of Sichuan Province. Bull. Chin. Mater. Med. 10: 398-400.
8. Wu, J. L. and Liu, X. C. 1989. Research on identification of medicinal "Tu-Sha-Ren". J. Chin. Med. Mater. 12: 19-24.
9. Chen, X., Huang, L. H., Long, C. C. and Mo, F. 2004. To distinguish *Alpinia zerumbet* from other Fructus Amomi. Guizhou Sci. 22: 53-55.
10. Lai, M. X., Chen, S. W., Fang, D. and Qin, D. H. 1989. Morphologic studies on the adulterant Sharen in *Alpinia*. China J. Chin. Mat. Med. 14: 267-270.
11. Lai, M. X., Chen, S. W., Fang, D. and Qin, D. H. 1989. Studies on the histology and essential oils of the adulterant Sharen in *Alpinia*. China J. Chin. Mat. Med. 14: 394-397.
12. Zhang, Y. B., Shaw, P. C., Sze, C. W., Wang, Z. T. and Tong, Y. 2007. Molecular authentication of Chinese Herbal Materials. J. Food Drug Anal. 15: 1-9.
13. Shaw, P. C., Ngan, F. N., But, P. H. and Wang, J. Molecular markers in Chinese medicinal materials. In "Authentication of Chinese Medicinal Materials by DNA Technology". 1st ed. pp. 1-24. Shaw, P. C., Wang, J. and But, P. H. eds. World Scientific Press. Singapore.
14. Wang, P. X., Huang, F., Zhou, L., Cao, L. Y., Liang, R. Y., Xu, H. H. and Liu, J. M. 2000. Analysis of *Amomum villosum* species and some adulterants of Zingiberaceae by RAPD. J. Chin. Med. Mater. 23: 71-74.
15. Xu, J. Y. and Ding, P. 2005. Analysis on genuineness of *Amomum villosum* Lour. From different habitats by RAPD. Tradit. Chin. Drug Res. Clin. Pharmacol. 16: 194-196.
16. Pan, H. X., Huang, F., Wang, P. X., Zhou, L., Cao, L. Y. and Liang, R. Y. 2001. Identification of *Amomum villosum*, *Amomum villosum* var. *xanthioides* and *Amomum longiligulare* on ITS1 sequence. J. Chin. Med. Mater. 24: 481-483.
17. Zhou, L., Wang, P. X., Huang, F., Cao, L. Y. and Liang, R. Y. 2002. ITS sequence analysis of *Amomum villosum*. Chin. Tradit. Herb. Drugs 33: 72-75.
18. Rogers, S. O. and Bendich, A. J. 1985. Extraction of DNA from milligram amounts of fresh, herbarium and mummified plant tissues. Plant Mol. Bio. 5: 69-76.
19. Douzery, E. J., Pridgeon, A. M., Kores, P., Linder, H. P., Kurzweil, H. and Chase, M. W. 1999. Molecular phylogenetics of *Diseae* (Orchidaceae): a contribution from nuclear ribosomal ITS sequences. Am. J. Bot. 86: 887-899.