



2009

Effect of chloramphenicol on a bioassay response for the detection of tetracycline residues in milk

Follow this and additional works at: <https://www.jfda-online.com/journal>

Recommended Citation

Nagel, O.; Zapata, M. de la L.; Basílico, J.C.; Bertero, J.; Molina, M.P.; and Althaus, R. (2009) "Effect of chloramphenicol on a bioassay response for the detection of tetracycline residues in milk," *Journal of Food and Drug Analysis*: Vol. 17 : Iss. 1 , Article 4.

Available at: <https://doi.org/10.38212/2224-6614.2303>

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

Effect of Chloramphenicol on a Bioassay Response for the Detection of Tetracycline Residues in Milk

ORLANDO NAGEL¹, MARÍA DE LA LUZ ZAPATA², JUAN CARLOS BASÍLICO², JORGE BERTERO¹,
MARÍA PILAR MOLINA³ AND RAFAEL ALTHAUS^{1*}

¹ Cátedra de Biofísica, Departamento de Ciencias Básicas,
Universidad Nacional del Litoral, R.P.L. Kreder 2805, 3080 Esperanza, República Argentina.

² Cátedra de Microbiología, Facultad de Ingeniería Química,
Universidad Nacional del Litoral, Santiago del Estero 2826, 3000 Santa Fe, República Argentina.

³ Departamento de Ciencia Animal, Universidad Politécnica de Valencia, Camino de Vera, Valencia, España.

(Received: February 4, 2008; Accepted: October 9, 2008)

ABSTRACT

The tetracyclines are commonly used in veterinary medicine, and yet the residues are not always detected by microbiological inhibitor tests using *Geobacillus stearothermophilus* subsp. *calidolactis* C-953 at their Maximum Residue Limit levels. In order to improve the sensitivity of these methods, a bioassay was evaluated to study the effect produced by the incorporation of different chloramphenicol (CAP) concentrations in the culture medium. The specificity and detection limits of six tetracyclines in milk were determined. As the levels of CAP increased, a decrease in specificity, from 97.9% (for 0, 50, 200 and 400 µg/kg of CAP) to 88.0% (for 600 µg/kg of CAP) was observed. The logistic regression model indicates a significant effect of the CAP concentration. However, the tetracycline-CAP interaction was not significant, and thus, a synergetic effect can not be considered between the two antimicrobials, with only a simple sum of their antimicrobial effects. When the CAP concentration is increased from 0 to 400 µg/kg, the detection limits of chlortetracycline (590 µg/kg-316 µg/kg), doxycycline (115 µg/kg-62 µg/kg), meclocycline (105 µg/kg-52 µg/kg), oxytetracycline (446 µg/kg-273 µg/kg), rolitetracycline (191 µg/kg-134 µg/kg) and tetracycline (302 µg/kg-158 µg/kg) are observed to decrease.

Key words: tetracycline, chloramphenicol, milk, bioassay, microbiological inhibitor test

INTRODUCTION

Tetracyclines (TCs) are broad-spectrum agents, exhibiting activity against infections caused by both Gram-positive and Gram-negative bacteria, as well as chlamydia, mycoplasmas, rickettsiae, and protozoan parasites.

Oxytetracycline (OTC), chlortetracycline (CTC) and tetracycline (TC) are widely used in the field of veterinary medicine for treating mastitis and metritis in cows. OTC and TC are also frequently added at subtherapeutic concentrations to cattle feeds, primarily, for collective prophylaxis. Other TCs, such as doxycycline (DC), meclocycline (MC) and rolitetracycline (RTC) are not currently used, but may be available commercially⁽¹⁾.

Either the inappropriate use of these drugs or the inadequate withdrawal period of treated animals can reveal TCs residues⁽¹⁻²⁾. One of the most worrying problems associated with TCs residues is the idiosyncratic

reactions, especially in hypersensitive consumers⁽³⁻⁵⁾. Like other antibiotics, TCs may be agents for the selection of antibiotic-resistant organisms⁽⁶⁾. In addition, some residues can interfere with the starter cultures used in processed milk products⁽⁷⁾. Thus, Mäyrä-Mäkinen⁽⁸⁾ has reported that TC of 0.3 µg/mL and 0.7 µg/mL modified the organoleptic properties of Edam and Emmenthal cheeses, respectively.

To prevent these problems, the European Union (EU)⁽⁹⁾ and the Codex Alimentarius⁽¹⁰⁾ have set a Maximum Residue Limit (MRL) for OTC, CTC and TC (100 µg/kg) in foods of animal origin such as meat, milk, and eggs.

An analytical method for the routine monitoring of TCs residues in milk must be rapid, precise and economical in cost and time⁽¹¹⁾. The bioassay techniques are widely used as screening methods⁽¹²⁾, since they are capable of analyzing a large number of samples. In later stages, confirmation and quantification methods are required⁽¹³⁻¹⁶⁾.

A wide variety of screening methods elaborated by

* Author for correspondence. Tel: +00-54-3496-420639 ext.131 ;
E-mail: ralthaus@fcv.unl.edu.ar

different laboratories is available on the market⁽¹⁷⁻¹⁹⁾. Microbiological inhibition methods using *Geobacillus stearothermophilus* subsp. *calidolactis* C-953 as a biological sensor permit relatively quick responses are significantly remarkable since responses are obtained (about 2.5 to 3.5 hr) due to high temperature incubation (64°C). These microbiological inhibitor tests show tetracycline detection limits above the MRLs, which does not make them suitable for the analysis of this antibiotic group⁽²⁰⁻²²⁾.

In order to improve the sensitivity of the microbiological methods for the detection of TCs, some authors recommend the use of chelating agents in the culture medium such as glycol-bis (2-aminoethylether) N,N,N',N'-tetraacetic acid (EGTA)⁽²³⁾ or salts of oxalic acid⁽²⁴⁻²⁵⁾, as these substances capture the bivalent ions (Ca⁺², Mg⁺²) present in milk. The TCs have a high affinity for these bivalent ions and chelating agent incorporation improve the availability of TCs, thus reducing the detection limits of these microbiological methods.

Adding CAP to the culture medium is another way of improving the sensitivity, since the combined action with TCs produces a more pronounced inhibition of protein synthesis in bacteria. In fact, the TCs interfere with the binding of aa- tRNA with A site of 30S ribosomal subunit⁽²⁶⁾, which inhibits the elongation of the peptidic chain. In a similar way the CAP interfere tRNA binding to 50S ribosomal subunit⁽²⁷⁾.

Nouws *et al.*⁽²⁸⁾ incorporated 100 µg/kg CAP in order to improve the TCs detection limits on the plate in which *Bacillus cereus* is used. Unlike *G. stearothermophilus*, when *B. cereus* is used, the response time is extended to 18-24 hours. Thus, this bioassay is not practical to be implemented as a routine method.

Nevertheless, there are no available studies to investigate the effect of CAP on the sensitivity of *G. stearothermophilus* to TCs in a microtiter-plate test with dichotomous response. Due to this, the aim of this work was to evaluate the incorporation of CAP into the culture medium of a bioassay to determine its antimicrobial action.

MATERIALS AND METHODS

I. Preparation of the microplates

The culture medium, made of casein peptone (5 g/L), yeast extract (2.5 g/L), glucose (1 g/L) and agar (15 g/L), was sterilized to 121°C for 15 min, cooled to 50 ± 1°C and the pH was adjusted to 7.0±0.1.

The culture medium was inoculated with a spore suspension of *G. stearothermophilus* subsp. *calidolactis* C-953 (1 × 10⁷ spores/mL, Merck®, Ref. 1.11499) and the bromocresol purple indicator (0.05 mg/mL) was added. The culture medium was divided into aliquots that were fortified with the CAP solution to obtain the following concentrations:

Specificity of bioassay: CAP = 0, 50, 200, 400 and

600 µg/kg. There were four plates for each level of CAP (20 microplates).

Detection limits: CAP = 0, 50, 200 and 400 µg/kg. There were twelve plates for each level of CAP (48 microplates).

For each aliquot, every well of the ELISA microplates was filled with 100 µL of culture medium, sealed with adhesive bands and conserved at 4°C until use.

II. Specificity of bioassay

The animals came from cattle herds in Las Colonias (Santa Fe, Argentina). For the specificity study, 192 milk samples taken from drug-free animals were analyzed.

The milk samples selected had a chemical composition and pH values considered as habitual for bovine milk, low somatic cell counts (SCC < 400000 cells/mL) and a bacterial count acceptable for cow milk (CFU < 100000 cfu/mL).

The milk samples were analysed in duplicate on ELISA microplates fortified with the five previously detailed CAP levels (CAP= 0, 50, 200, 400 and 600 µg/kg).

For this purpose, 50 µL of milk samples were added in each individual well of the microplates. It was left at 4°C for one hour to allow the diffusion of the natural inhibitors of the milk to take place. Next, each microplate was washed several times with distilled water, floated in a water bath and then incubated at 64 ± 1°C for 3.5 hours.

Each sample of milk was also analysed in duplicate using the Copan® test (CH ATK) microplate P&S (Chr Hansen, Hoersholm, Denmark), so that their results can be compared with the specificity of the bioassay fortified with different levels of CAP.

A visual interpretation was carried out by 3 qualified people and evaluated as "negative" or "positive". "Doubtful" qualifications were considered "positive"⁽²⁹⁾.

The specificity of the bioassay was calculated as follows:

$$\text{Specificity} = \frac{\text{negative}}{\text{total samples}}$$

III. Detection Limits of Bioassay

Milk samples were fortified with six TCs: chlortetracycline (CTC), doxycycline (DC), meclocyline (MC), oxytetracycline (OTC), rolitetracycline (RTC) and tetracycline (TC), as detailed in Table 1.

These drugs were stored and handled according to the manufacturers' instructions before use. All dilutions were prepared in 10 mL volumetric flasks at the moment when the analyses were carried out, in order to avoid any possible inconvenience due to solution instability.

The antimicrobial solutions were prepared in one step only from the respective stock solution using antimicrobial-free milk (Copan® test microplate P&S, Chr Hansen, Hoersholm, Denmark). The final drug concentrations in milk (µg/kg) were reached after serial dilutions, in such a way that the volume of the antimicrobial

Table 1. Tetracycline concentrations used in the experiment

TCs	Product number	Concentrations ($\mu\text{g}/\text{kg}$)
CTC	Sigma C-4881	0, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 800
DC	Sigma D-9891	0, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140
MC	Sigma M-1388	0, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140
OTC	Sigma O-5750	0, 25, 50, 75, 100, 125, 150, 175, 200, 300, 400, 500
RTC	Sigma R-2253	0, 20, 40, 60, 80, 100, 120, 140, 160, 180, 200, 220
TC	Sigma T-3258	0, 25, 50, 75, 100, 125, 150, 175, 200, 300, 400, 500

CTC: Chlortetracycline, DC: Doxycycline, MC: Meclocyline, OTC: Oxytetracycline, RTC: Rolitetracycline, TC: Tetracycline.

agent solution did not exceed 1% of the volume of the final solution to be analysed⁽³⁰⁻³¹⁾.

For each tetracycline and CAP level, 16 replicates from each of the twelve concentrations detailed in Table 1 were analysed (2 microplates for each level of CAP, 8 microplates for each tetracycline).

A volume of 50 μL of each solution was used for the bioassay. The temperatures, diffusion and incubation time detailed in the specificity study were used. The visual interpretation was also carried out by 3 qualified people, and the test results were evaluated as “negative” or “positive”.

IV. Statistical Analysis

As the visual evaluation of the bioassay is an ordinal variable at two dichotomous levels (“negative” and “positive”), it is appropriate to use a logistic model to treat the data⁽³²⁾.

To analyze the results, the stepwise procedure was applied to the logistic regression option of SAS⁽³³⁾, taking the model of those variables that presented a value of Chi-squared ≥ 3.94 to be relevant. The logistic regression model was the following:

$$L_{ijk} = \text{logit} [P_{ijk}] = \beta_0 + \beta_1 [T]_i + \beta_2 [CAP]_j + \beta_{12} ([T][CAP])_{ij} + \varepsilon_{ijk}$$

Where: L_{ijk} = the dependent or response variable of the linear logistic model; $[P_{ijk}] = [P_p / (1-P_p)]$ or the ratio of the probability of “positive” response/the probability of “negative” response; $[T]_i$ = effect of tetracycline concentration ($i = 1, 2, \dots, 12$ levels, Table 1), $[CAP]_j$ = effect of CAP concentrations ($j = 0, 50, 200$ or $400 \mu\text{g}/\text{kg}$), $([T][CAP])_{ij}$ = effect of interaction between tetracycline and CAP concentrations; $\beta_0, \beta_1, \beta_2,$ and β_{12} = coefficients estimated for terms of intercept, tetracycline, CAP and interaction between tetracycline and CAP, respectively; and ε_{ijk} = residual error.

The concordance coefficient⁽³³⁾ was applied as the rank correlation between the observed responses and

predicted probabilities.

The detection limit of the visual interpretation of the bioassay was estimated as the concentration that produces 95% of the “positive” results^(29,31).

RESULTS AND DISCUSSION

I. Specificity of Bioassay

The results obtained from the analysis of 192 antibiotic-free milk samples are shown in Table 2. It is shown that the addition of $400 \mu\text{g}/\text{kg}$ of CAP in the culture medium does not produce an increase in the frequency of positive results (specificity = 97.9%), obtaining the same specificity as the CH-ATK[®] Test.

However, if compared with the microplates prepared with $400 \mu\text{g}/\text{kg}$ (23 vs. 4), the incorporation of $600 \mu\text{g}/\text{kg}$ of CAP does produce an increase in the frequency of positive results to the bioassay accompanied by an appreciable decrease in their specificity (88.0%). For this reason, the assay of the detection limits of TCs in milk samples was carried out with $400 \mu\text{g}/\text{kg}$ of CAP.

It should be noted that the specificity (97.9%) of

Table 2. Effect of the levels of CAP on the specificity of the bioassay

CAP	Sample number	Negative	Positive	Specificity (%)
CH-ATK [®] Test	192	188	4	97.9
0 $\mu\text{g}/\text{kg}$	192	188	4	97.9
50 $\mu\text{g}/\text{kg}$	192	188	4	97.9
200 $\mu\text{g}/\text{kg}$	192	188	4	97.9
400 $\mu\text{g}/\text{kg}$	192	188	4	97.9
600 $\mu\text{g}/\text{kg}$	192	169	23	88.0

Specificity: “negative/total samples”.

Table 3. Significance of the effects calculated by means of the logistic regression model

Tetracyclines	Factors	" χ^2 " value	"p" value
Chlortetracycline	[CTC]	148.3518	0.0001
	[CAP]	97.3408	0.0001
	[CTC]*[CAP]	0.0531	0.8177
Doxycycline	[DC]	104.8903	0.0001
	[CAP]	59.4681	0.0001
	[DC]*[CAP]	0.9314	0.3345
Meclocycline	[MC]	94.4473	0.0001
	[CAP]	58.6459	0.0001
	[MC]*[CAP]	0.9857	0.3208
Oxytetracycline	[OTC]	149.6728	0.0001
	[CAP]	75.8476	0.0001
	[OTC]*[CAP]	0.2845	0.5937
Rolitetracycline	[RTC]	149.0186	0.0001
	[CAP]	63.3352	0.0001
	[RTC]*[CAP]	0.0025	0.9604
Tetracycline	[TC]	113.0115	0.0001
	[CAP]	100.0742	0.0001
	[TC]*[CAP]	0.0637	0.8008

CTC: Chlortetracycline, DC: Doxycycline, MC: Meclocycline, OTC: Oxytetracycline, RTC: Rolitetracycline, TC: Tetracycline, CAP: Chloramphenicol.

microplates fortified with 400 $\mu\text{g}/\text{kg}$ of CAP was similar to the values of 98%⁽³⁴⁾ and 95%⁽³⁵⁾ reached when the Delvotest[®] test in cow milk samples.

II. Detection Limits of Bioassay

Table 3 summarises the results by applying the logistic regression model. It is shown that the levels of CAP in the culture medium significantly affected ($p < 0.0001$) the response of the bioassay for the six TCs analysed. The interaction of the different TCs with CAP, however, was not significant ($p > 0.05$) in the six studied cases.

This fact indicates that CAP produces an antimicrobial effect in the bioassay, but there is no synergetic action between the two antimicrobial agents, although both antimicrobials participate at tRNA level and impede the synthesis of proteins.

In fact, statistical model points to a simple additive antimicrobial effects of CAP and TCs, as opposed to a synergetic effect, such as, sulphonamides and antifolates (e.g. trimetoprim, tetroxoprim, pirimetamine).

The coefficients obtained for six TCs by means of a logistic regression model are shown in Table 4. The

values of " β_1 " coefficients indicate the increase in the frequency of positive results as the TC concentration increases in the milk samples. It can be observed that DC ($\beta_1 = 0.0960$) and MC ($\beta_1 = 0.1057$) possess higher values of this coefficient compared with CTC ($\beta_1 = 0.0216$) and OTC ($\beta_1 = 0.0249$). The higher " β_1 " coefficient values indicate a smaller increment in their concentrations to produce 100% positive results, while TCs have lower values of the " β_1 " coefficients, thereby requiring greater concentrations in order to reach 100% positive results. Similarly, the " β_2 " coefficients represent the antimicrobial action caused by the CAP in the culture medium. The fact that the values of these coefficients (Table 4) for the six analysed TCs fell between $\beta_2 = 0.0108$ (OTC) and $\beta_2 = 0.0142$ (MC), indicating that the CAP's antimicrobial action performed in a similar way.

It may be observed in Table 4 that the concordance coefficients were high, ranging between 97.3% (RTC) and 99.0% (TC), which demonstrate the close fit of the model.

Figure 1 shows the effect of the concentrations of TCs and CAP on the relative frequency of positive results as well as the curves constructed by means of the logistic regression model (" β_0 ", " β_1 " and " β_2 " coefficients, Table 4). An increase in CAP concentration produces a displacement of the dose-response curve to lower concentrations, improving the sensitivity of the bioassay.

Table 4. Summary of logistic regression model coefficients

Tetracyclines	β_0	β_1	β_2	β_{12}	C
Chlortetracycline	-9.8056	0.0216	0.0148	-	98.2 %
Doxycycline	-8.1365	0.0960	0.0128	-	98.3 %
Meclocycline	-8.1833	0.1057	0.0142	-	98.5 %
Oxytetracycline	-8.1687	0.0249	0.0108	-	98.6 %
Rolitetracycline	-10.1647	0.0685	0.00983	-	97.3 %
Tetracycline	-10.2874	0.0438	0.0158	-	99.0 %

C: concordance coefficients.

Table 5. Effect of the concentration of CAP on the detection limits ($\mu\text{g}/\text{kg}$) of tetracyclines in milk

Tetracyclines	Concentration CAP (mg/kg)				MRLs*
	0	50	200	400	
Chlortetracycline	590	556	453	316	100
Doxycycline	115	109	89	62	-
Meclocycline	105	99	78	52	-
Oxytetracycline	446	425	360	273	100
Rolitetracycline	191	184	163	134	-
Tetracycline	302	284	230	158	100

*MRLs ($\mu\text{g}/\text{kg}$), EU maximum residue limits, CAP: Chloramphenicol.




Figure 1. Tetracycline dose-response curves for different chloramphenicol concentrations (\square CAP: 0 $\mu\text{g/mL}$, \diamond CAP: 50 $\mu\text{g/mL}$, Δ CAP: 200 $\mu\text{g/mL}$, \times CAP: 400 $\mu\text{g/mL}$).

The detection limits of the bioassay calculated by means of the logistic model and the 95% relative frequency of positive results⁽²¹⁾ are summarised in Table 5, along with the values of the Maximum Residue Limits set out by the European Union (EU-MRLs). The incorporation of CAP into the culture medium decrease the detection limits of six TCs, although this decrease does not reach the values of EU-LMRs.

The detection limits of OTC (273 $\mu\text{g/kg}$) and TC (158 $\mu\text{g/kg}$) calculated in the bioassay containing 400 $\mu\text{g/kg}$ of CAP (Table 5) are lower than the 400 $\mu\text{g/kg}$ ⁽³⁶⁾ or 500 $\mu\text{g/kg}$ ⁽³⁷⁾ of OTC and 600 $\mu\text{g/kg}$ ⁽³⁸⁾ of TC determined in the Delvotest[®] SP microbiological inhibition test using *G. stearothersophilus*.

The BRT[®] AiM test, which uses a redox indicator (Brilliant Black) in bioassay response, presents very high detection limits. Frank⁽³⁹⁾ obtained values of about 5000

$\mu\text{g/kg}$ for OTC and TC residues in cow milk samples.

It should be mentioned that TCs possess a tendency to form chelate complexes with ions of bivalent metals⁽⁴⁰⁾. However, TCs can be displaced from these complexes by the addition of stronger chelating agents such as EGTA⁽²³⁾. In order to improve the sensitivity of *G. stearothersophilus* to TCs, further studies should be carried out by means of the incorporation of CAP and chelating agents of calcium so that the detection limits could approach those of the Maximum Residue Limits.

CONCLUSIONS

The incorporation of 400 $\mu\text{g/kg}$ of CAP into the culture medium of a bioassay that contains *G. stearothersophilus* diminishes the detection limits of the TCs in

milk without affecting the specificity. Also, such CAP concentrations allow us to achieve detection limits which are near those of the Maximum Residue Limits.

This decrease in the detection limits is due to a sum of the antimicrobial effects of the CAP and TCs, rather than a synergetic effect of the two antimicrobial agents.

In order to approach the Maximum Residue Limits of TCs in milk, further studies should be carried out that contemplate the combined incorporation of CAP and chelating agents into the culture medium of the bioassay.

ACKNOWLEDGEMENTS

This investigation was carried out as part of the CAI+D'05/07 Project (N° 033-213 Res. H.C.D.N° 302), supported by financial assistance from the Universidad Nacional del Litoral (Santa Fe, República Argentina). The authors also thank the Proyecto Federal de Innovación Productiva (PFIP 2006, Res. 1888/05 del 22/12/05).

REFERENCES

- Carson, M. C., Ngoh, M. A. and Hadley, S. W. 1998. Confirmation of multiple tetracycline residues in milk and oxytetracycline in shrimp by liquid chromatography-particle beam mass spectrometry. *J. Chromatogr.* 712: 113-128.
- Samanidou, V., Nikolaidou, K. and Papadoyannis, I. 2005. Development and validation of an HPLC confirmatory method for the determination of tetracycline antibiotics residues in bovine muscle according to the European Union regulation 2002/657/EC. *J. Sep. Sci.* 28: 2247-2258.
- Huber, W. G. 1986. Allergenicity of Antibacterial Drug Residues. In "Drug Residues in Animal". pp.33-50. Academic Press, Inc. New York, U.S.A.
- Collins-Thompson, D. L., Wood, D. S. and Thomson, I. Q. 1988. Detection of antibiotic residues in consumer milk supplies in North America using the Charm II test procedure. *J. Food Prot.* 51: 632-633.
- Katz, S. E. and Brady, M. S. 1988. Antibiotic Residues in Food and Their Significance. In "Antimicrobials in Foods". Davidson P., Branen, A. L. Ed. pp. 571-595. Merce Dekker. New York, U.S.A.
- Brady, M. S., Strobel, R. J. and Katz, S. E. 1988. Vitro Analytical System for Determining the Ability of Antibiotic at Residue Levels to Select for Resistance in Bacteria. *J. A. O. A. C.* 71: 295-298.
- Claypool, R. H. 1984. Concerns and Problem Cheese Industry for Milk Quality and Procurement-Today and Future. *J. Dairy Sci.* 67: 2091-2094.
- Mäyrä-Mäkinen, A. 1995. Technological significance of residues for the dairy industry. In "Residues of Antimicrobial Drugs and Other Inhibitors in Milk". IDF S. I. N° 9505. International Dairy Federation. pp. 136-143. Brussels, Belgium.
- Commission of the European Communities. 1991. Establishment by the European Community of Maximum Residue Limits (MRLs) for residues of veterinary medical products in foodstuffs of animal origin. The rules governing medical products in European Community VI. ECSC-EEC-EAEC. Brussels, Belgium.
- Codex Alimentarius. 1996. Residuos de medicamentos veterinarios en los alimentos. Codex Alimentarius, 3. Programa conjunto FAO/OMS sobre Normas Alimentarias, FAO. Roma, Italy.
- Furusawa, N. 2003. Isolation of tetracyclines in milk using a solid-phase extracting column and water eluent. *Talanta* 59: 155-159.
- Honkanen-Buzalski, T. and Reybroeck, W. 1995. Antimicrobials. In "Residues of antimicrobial drugs and other inhibitors in milk". IDF Special Issue N° 9505. International Dairy Federation. pp. 26-35. Kiel, Germany.
- Blanchflower, W., McCracken, R. and Desmond, R. 1989. Determination of chlortetracycline residues in tissues using high-performance liquid chromatography with fluorescence detection. *Analyst* 114: 421-423.
- Cinquina, A., Longo, F., Anastasi, G., Giannetti, L. and Cozzani, R. 2003. Validation of a high-performance liquid chromatography method for the determination of oxytetracycline, tetracycline, chlortetracycline and doxycycline in bovine milk and muscle. *J. Chromatogr. A* 987: 227-233.
- Zhao, F., Zhang, X. and Gan, Y. 2004. Determination of tetracyclines in ovine milk by high-performance liquid chromatography with a coulometric electrode array system. *J. Chromatogr. A* 1055: 109-114.
- Biswas, A., Rao, G., Kondaiah, N., Anjaneyulu, A., Mendiratta, S., Prasad, R. and Malik, J. 2007. A simple multi-residue method for determination of oxytetracycline, tetracycline and chlortetracycline in export buffalo meat by HPLC-photodiode array detector. *J. Food Drug Anal.* 15: 278-284.
- International Dairy Federation (IDF). 1991. Detection and confirmation of inhibitors in milk and milk products. IDF Bull. N° 258. International Dairy Federation. Bruselas, Belgium.
- Zorraquino, M., Berruga, M. and Solaz, T. 2003. Análisis de las técnicas de detección de residuos de medicamentos veterinarios en la leche cruda y sistema de control. Informe Técnico. Federación Industrias Lácteas Españolas (INLAC). Madrid.
- Diserens, J. 2005. Review of available screening tests for the detection/determination of veterinary drug residues in milk. International Dairy Federation JAT on Veterinary Residues meeting, Item 5. Parma, Italy.
- Van Os, J. and Beukers, R. 1980. A multitest system for detection of antibiotics in milk. *J. Food Prot.* 43: 510-511.
- Luitz, M., Suhren, G. and Heeschen, W. 1996. Interactions in milk consequences for the detection by a

- microbial inhibitor test. *Milchwissenschaft* 51: 390-392.
22. Abete, M., Di Falco, M., Genta, E. and Squadrone, S. 1997. Contronto tra i metodi microbiologici e i metodi immunoenzimatici per el rilevamento delle tetraciline nel latte. *Il Latte*. 22: 74-77.
 23. Adriany, A., Märtlbauer, E. and Zaadhof, K. 1995. A modified brilliant black-reduction test (BRT) with improved sensitivity for tetracyclines and sulfonamides. *Proceedings of the IDF-Symposium Residues of antimicrobial drugs and other inhibitors in milk*. 28-31 August 1995. pp. 172-174. Kiel, Germany.
 24. Langeveld, P. 2005. Improved method for the determination of the presence of an antibiotic in fluid. World Intellectual Property Organization. Patent n° WO 2005/005656 A1 of 20 Jan. 2005.
 25. Langeveld, P., Beukers, R., Bommele, M. and Stark, J. 2005. Rapid microbiological test for the detection of antibacterial compounds. United States Patent. Patent n° US 6.867.015 B1 of 15 of Mar. 2005.
 26. Triton, T. 1977. Ribosome-Tetracycline Interactions. *Biochemistry* 16: 4133-4138.
 27. Lessard, J. and Pestka, S. 1972. Studies on the formation of transfer ribonucleic acid-ribosome complexes. XXIII. Chloramphenicol, aminoacyl-oligonucleotides and *Escherichia coli* ribosomes. *J. Biol. Chem.* 247: 6909-6912.
 28. Nouws, J., Van Egmond, H., Sholders, I., Loeffen, G., Schouten, J. and Stegeman, H. 1999. A microbiological assay system for assessment of raw milk exceeding EU maximum residue level. *Int. Dairy J.* 9: 85-90.
 29. Suhren, G., Reichmuth, J. and Walte, H. 1996. Detection of β -lactam antibiotics in milk by the Penzym-test. *Milchwissenschaft* 51: 269-273.
 30. Reichmuth, J. Suhren, G. and Beukers, R. 1997. Evaluation of microbial inhibitor test-the IDF approach. *Milchwissenschaft* 52: 691-695.
 31. International Dairy Federation (IDF). 1999. Guidance for the standardized evaluation of microbial inhibitor test. IDF Standard N° 183. International Dairy Federation. Brussels, Belgium.
 32. Agresti, A. 1990. *Categorical Data Analysis*. Ed. John Wiley and Sons. New York, U.K.
 33. SAS® Institute Inc., 2001. *SAS Users guide: statistics version 9.1*. Cary, NC.
 34. Sischo, W. and Burns, C. 1993. Field trial of four cowside antibiotic residue screening test. *J. Am. Vet. Med. Assoc.* 202: 1249-1254.
 35. Charm, S. and Zomer, E. 1995. The evolution and direction of rapid detection/Identification of antimicrobial drug residues. In "Residues of antimicrobial drugs and other inhibitors in milk". IDF Special Issue N° 9505. *Int. Dairy Fed.* pp. 224-233. Brussels, Belgium.
 36. Zaadhof, H., Mältlbauer, E., Vormeister, A. and Schweizer, L. 1997. Zur eignung kommerzieller mikrobiologischer hemmstofftests als suchverfahren auf das vorhandensein von antiinfektiva in milch und erzeugnissen auf milchbasis. *Archiv für Lebensmittelhygiene*. 48: 132-144.
 37. Luitz, M. and Suhren G. 1996. Nachweis von tetracyclinen in milch mit verschiedenen modifikationen des brillant-schwarzreduktionstests und dem Delvotest SP. DMZ, Lebensmittelindustrie und Milch-wirtschaft. 117: 1010-1015.
 38. Senyk, G., Davidson, J., Brown, J., Hallstead, E. and Sherbon, J. 1990. Comparison of rapid test used to detect antibiotic in milk. *J. Food Prot.* 53: 158-164.
 39. Frank, S. 1995. Bestimmung der Nachweisgrenzen ausgewählter Antibiotika und Sulfonamide im Brillantschwarz-Reduktionstest. *Diss. Universität Diplomarbeit Thier. Triesdorf. Germany.*
 40. Rogalsky, W. 1985. Chemical modification of the tetracyclines. In "The tetracyclines" Hlavka, J., Boothe, J., Springer, E. Ed. pp. 114-129. Berlin, Germany.