



2008

Factors affecting PGE2 production in seaweed *Gracilaria tenuistipitata*

Follow this and additional works at: <https://www.jfda-online.com/journal>

Recommended Citation

Hsu, B.-Y.; Tsao, C.-Y.; Chiou, T.-K.; and Hwang, D.-F. (2008) "Factors affecting PGE2 production in seaweed *Gracilaria tenuistipitata*," *Journal of Food and Drug Analysis*: Vol. 16 : Iss. 3 , Article 11.
Available at: <https://doi.org/10.38212/2224-6614.2358>

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

Factors Affecting PGE₂ Production in Seaweed *Gracilaria tenuistipitata*

BO-YANG HSU, CHING-YU TSAO, TZE-KUEI CHIOU AND DENG-FWU HWANG*

Department of Food Science, National Taiwan Ocean University, 2 Pei-Ning Rd., Keelung City 202, Taiwan (R.O.C)

(Received: March 15, 2007; Accepted: October 23, 2007)

ABSTRACT

Gracilaria tenuistipitata is an edible red alga and can be utilized as feed for small abalone and foodstuff (ex. agar extract). It is the major species of *Gracilaria* that is commonly cultivated and consumed in Taiwan. Effects of environmental factors on prostaglandin E₂ (PGE₂) production in the seaweed *G. tenuistipitata* were investigated in this study. PGE₂ amount increased nearly 2-fold and 47% when the seaweed was exposed to two ways of fluctuation of seawater salinity from 2‰ to 1‰ and 3‰ for 12 hr, respectively. The stresses of low temperature and irradiance promoted the seaweed to produce PGE₂. When the seaweeds were exposed to air for 2 hr and 4 hr, the PGE₂ levels were elevated by 25% and 31%, respectively. When the seaweed was soaked in freshwater, PGE₂ amount gradually decreased. Cu²⁺ and Zn²⁺ inhibited PGE₂ production at the concentrations of 3 and 50 mg/L, respectively. Mg²⁺ slightly inhibited PGE₂ production about 20% at 2,025 mg/L, while Ca²⁺ boosted PGE₂ production about 59% at 600 mg/L. Hence, environmental factors significantly affect the PGE₂ production in the seaweed *G. tenuistipitata*. Prostaglandins might be associated with processes permitting the algae tissue to survive in unfavorable conditions. To lower the amount of PGE₂ in the seaweed and ensure the food safety, it is suggested to minimize the variation of growth condition to avoid PGE₂ increase in the seaweed during cultivation period. In addition, consumers should avoid eating raw seaweed since PGE₂ could be destroyed by heating.

Key words: prostaglandin, arachidonic acid, *Gracilaria tenuistipitata*, environmental factors

INTRODUCTION

Gracilaria sp. is one kind of edible red algae (Japanese name, "ogonori"). They are generally utilized in foodstuff, food additive and cosmetic. People are accustomed to take raw fish with *Gracilaria* sp. in Japan^(1,2). Raw fish and red algae are rich in polyunsaturated fatty acids (PUFA), mainly arachidonic acid (AA) and eicosapentaenoic acids which are precursors of prostaglandins (PGs)^(3,4). However, some poisoning cases occurred due to ingesting seaweed *G. verrucosa*, known as ogonori, or similar species in Japan⁽¹⁻³⁾. Those victims ate raw fish and raw seaweed. The common symptoms in the patients are nausea, vomiting and diarrhea appearing 30-60 min after ingestion. In extreme cases, they developed very low blood pressure, followed by shock and death. In addition, the symptoms in female were more serious than those in male. The possible reason of seaweed poison cases was that raw seaweed contained cyclooxygenase (COX) which could transform AA (precursor of prostaglandin E₂, PGE₂) or other highly unsaturated fatty acids to prostaglandins, especially PGE₂. The enzyme in the seaweed and/or the body tissues of the victim may be

acting on the highly unsaturated fatty acids in the raw fish or in the seaweed producing a great amount of PGE₂ at short time in the body⁽¹⁾. PGE₂ has many physiological effects on mammals including hyperthermia, hypotension, smooth muscle dilatation, gastric secretion inhibition, hyperalgesia, womb shrink and relating to cancer⁽⁵⁻¹⁰⁾. Noguchi *et al.* reported that 15.8-102 ppm of PGE₂ and small amounts of other PGs (A₁, E₁ and F_{2α}) could be detected from poison sample. They assumed that AA from raw tuna slices and ogonori were catalyzed by the COX in ogonori and victim to form PGs which caused the poisoning symptoms⁽¹⁾.

Prostaglandins were also found in different species of *Gracilaria*, such as *G. asiatica* and *G. lichenoids* containing PGE₂^(2,11). Some studies reported that prostaglandin amounts were various with seasons and species of *Gracilaria*. For example, PGE₂ amounts in *G. asiatica* were much higher in winter than in summer^(2,3). There are several factors that can influence on prostaglandin amount in the different seasons, including stage of life cycle, activity of enzymes of related prostaglandins synthesis, environmental conditions (seawater temperature, irradiance, contents of dissolved components, etc.) and others⁽³⁾. Sajiki reported PGE₂ amount in *G. asiatica* was much higher than that in *G. rhodocaudata*. There-

* Author for correspondence. Tel: +886-2-2462-2192 ext. 5103; Fax: +886-2-2462-6602; E-mail: dfhwang@mail.ntou.edu.tw

fore PGE₂ concentrations in ogonori may depend on species and environmental conditions⁽²⁾.

G. tenuistipitata is one of ogonori and the major cultivated and utilized species in Taiwan. As described above, PGE₂ was the possible causative agent for "ogonori poison". Therefore the present study was aimed to investigate effects of environmental factors on PGE₂ production in *G. tenuistipitata*. The PGE₂ amount in the seaweed *G. tenuistipitata* was investigated when it was exposed to different environments, including the stress of salinity, temperature, irradiance, metal ions, tide exposure, and freshwater exposure. The change of growing conditions may exert influence on the PGs production in *G. tenuistipitata*.

MATERIALS AND METHODS

I. Chemicals and Solvents

Prostaglandin E₂ and arachidonic acid sodium salt (90% of purity) were purchased from Sigma Chemical Co. (St. Louis, Mo, USA). Acetonitrile was from Mallinckrodt Baker, Inc. (NJ, USA). Methanol, ethyl acetate, citric acid, sodium citrate, CuCl₂, ZnCl₂, MnCl₂, CaCl₂ and MgCl₂ were from Panreac Co. (Barcelona, Spain).

II. Materials

Raw seaweed (*G. tenuistipitata*) samples were collected from a seaweed farm in Hukou, Yunlin County, Taiwan. These samples were collected all year long.

III. Standards

Standard stock solutions of PGE₂ were made in acetonitrile. Suitable amount of standard PGE₂ was dissolved in acetonitrile to make 1 mg/mL stock solution of standard. Various concentrations (1, 5, 10, 25, 50 and 100 µg/mL) of standard solutions were prepared by serial dilution of the stock solution (1 mg/mL) with acetonitrile. The standard solutions were stored at -18°C because PGE₂ is unstable at room temperature.

IV. Sample Preparation for Analysis

Ten grams wet raw seaweed sample with ~90% moisture content was finely sliced and ground in a mortar. The centrifugation was performed at 13,700 ×g, 0°C for 30 min and the supernatant was collected. To the supernatant was added 2 mg of arachidonic acid sodium salt and incubated at 37°C for 30 min. The pH was adjusted to 3-4 with 1.5 M citric acid. The mixture was incubated in boiling water for 10 min and then cooled on ice. The solution was purified by solid phase extraction (SPE). The C-18 SPE cartridge (Mallinckrodt Baker, Inc. NJ,

USA) was activated by rinsing with 5 mL methanol and then with 5 mL distilled water. The sample was loaded over the C18 SPE cartridge and the cartridge was rinsed with 5 mL distilled water. The cartridge was then eluted with 2 mL methanol followed by 5 mL ethyl acetate⁽¹²⁾. The ethyl acetate extracts were evaporated to dryness under reduced pressure and the residue was dissolved in 1 mL acetonitrile as test solution. Test solution (10 µL) was submitted to HPLC analysis.

V. HPLC Analysis

The PGE₂ contents were determined by using a Hitachi Liquid Chromatography system (Hitachi, Ltd, Tokyo) consisting of a Model L-6200 pump, a Rheodyne Model 7125 syringe loading sample injector, a Model L-7455 diode array detector set at 196 nm and a Model D-7000 Chromato-integrator. Analysis of PGE₂ was performed on a YMC-Pack ODS-AQ AQ 312 (150 × 6.0 mm I.D., S-5 µm, 120 Å) reversed phase column⁽¹⁾.

The mobile phase is composed of 17 mM phosphoric acid and acetonitrile gradient from 35% to 60% in 30 min. The flow rate was 1.0 mL/min⁽¹³⁾.

VI. Standard Curve and Sample Quantification

Standard curve was obtained in the range of 1~100 µg/mL for PGE₂. The curve was plotted by standard peak area versus standard concentration in µg/mL. Amount of PGE₂ in the sample was calculated out of the standard curve.

VII. Effect of Environmental Factors on PGE₂ Production

(I) Effect of Salinity on PGE₂ Production

The seaweed *G. tenuistipitata* was acclimated in seawater with salinity 2% at least 3 days, moved to salinity 1% and 3% seawater and then incubated for 12 hr. All samples were then submitted to the extraction and PGE₂ analysis by HPLC.

(II) Effect of Temperature on PGE₂ Production

The seaweed was acclimated in seawater (25°C) at least 3 days, moved to 4, 15 and 35°C seawater, and incubated for 12 hr. All samples were then submitted to the extraction and analysis.

(III) Effect of Irradiance on PGE₂ Production

The seaweed was acclimated in seawater with irradiance 330 µE m⁻² s⁻¹ for 3 days, moved to seawater with different irradiance including 0 and 640 µE m⁻² s⁻¹, and then incubated for 12 hr. Light meter was used to monitor the irradiance (Lutron LX-102) in the range of 0 to 1,000 µE m⁻² s⁻¹. All samples were submitted to the extraction and analysis.

(IV) Effect of Air Exposure on PGE₂ Production

The seaweed was acclimated in seawater at least 3 days and then exposed to air for 2 and 4 hr. All samples were submitted to the extraction and analysis.

(V) Effect of Freshwater Exposure on PGE₂ Production

The seaweed was acclimated in seawater at least 3 days and then soaked in freshwater for 2 and 4 hr. All samples were submitted to the extraction and analysis.

(VI) Effects of Metal Ions on PGE₂ Production

According to the standard amounts of various metal ions in ocean from Environmental Protection Administration, Executive Yuan, Taiwan, ROC, the standard levels of Zn²⁺, Cu²⁺ and Mn²⁺ are 0.5, 0.03 and 0.05 mg/L (control concentration), respectively. The concentrations of Zn²⁺, Cu²⁺ and Mn²⁺ in this study were 10 and 100 folds higher than the standards. The common levels of Ca²⁺ and Mg²⁺ in the ocean are 400 and 1350 mg/L (control concentration), respectively. The concentrations of Ca²⁺ and Mg²⁺ in this study were 50% higher than average concentration in seawater. The seaweed was acclimated in seawater at least 3 days and cultivated in seawater with respectively containing Zn²⁺, Cu²⁺, Mn²⁺, Ca²⁺, and Mg²⁺ for 12 hr. All samples were submitted to the extraction and analysis.

VIII. Statistical Analysis

All measurements were conducted in triplicate, and experimental results were evaluated by analysis of variance (ANOVA) (SAS Version 9.0, SAS Inst. Inc., Cary, NC, USA) and Duncan's multiple range tests was applied to compare the mean values at $P < 0.05$.

RESULTS AND DISCUSSION

The mobile phase of HPLC for PGE₂ was as follows: the ratio of acetonitrile and 17 mM phosphoric acid changed from 35/65 to 60/40 in 30 min. The flow rate was 1.0 mL/min⁽¹³⁾. Typical chromatograms are shown in Figure 1. The retention time of PGE₂ was 12.82 ± 0.20 min. The peak in the sample at retention time 21 min was suggested to be PGA₂ by comparing to previous datum⁽¹³⁾. However, the other two peaks in the sample at retention time 8 min and 15.5 min were unknown peaks.

The effect of salinity fluctuation on PGE₂ amount in *G. tenuistipitata* is shown in Figure 2. The optimal growth salinity of *G. tenuistipitata* is 1.5-2.5%. Therefore the seaweed is cultivated in 2% salinity seawater (control condition). When the seaweed was cultivated in 2% salinity seawater and then transferred to 1% and 3% salinity seawaters after 12 hr, PGE₂ amount in the




Figure 1. HPLC chromatograms of PGE₂ standard (A), along with sample (B). Acetonitrile concentration increased from 35 to 60% in 30 min and the flow rate was 1.0 mL/min.




Figure 2. PGE₂ amount of *Gracilaria tenuistipitata* in different salinities. The seaweed was cultivated in 2% salinity seawater and then transferred to salinity 1% and 3% for 12 hr, respectively. Values are mean ± S.D. (N = 3). a, b, c: Values with different superscripts are significantly different at $P < 0.05$.

seaweed increased, resulting in the highest production in 1% salinity seawater, followed by 3% salinity seawater and control (2% salinity seawater). PGE₂ amount increased nearly 2-folds and 47%, respectively, when the seaweed was transferred from 2% to 1% and 3% salinity seawaters. Dawes *et al.* reported low salinity (1.5%) increased the level of AA in *G. tikvahiae* sporelings⁽¹⁴⁾. The levels of PGE₂ and AA were increasing at the same time in lower salinity; therefore there could be positive relationship between AA and prostaglandins due to AA being a precursor of prostaglandins. The result indi-

cated that conversion of salinity caused the seaweed to respond to the stress from environmental change due to the role of PGE₂ in homeostasis of seaweed.

PGE₂ amounts in *G. tenuistipitata* were different when the seaweed was transferred to different temperatures of seawater after 12 hr. The optimal growth temperature of *G. tenuistipitata* is 20-25°C. Therefore seaweed is cultivated in 25°C (control condition). PGE₂ amount increased 49% and 76% when the seaweed was transferred from 25°C to 4 and 15°C, respectively after 12 hr as shown in Figure 3. In addition, PGE₂ amount reduced 45% when the seaweed was transferred from 25°C to 35°C after 12 hr. Imbs *et al.* reported that there were different prostaglandin contents in *G. verrucosa* during June-November. The levels of PGE₂ and PGF_{2α} did not show significant difference in June-September, but showed the highest amounts in November when water temperature is relatively low⁽³⁾. Sajiki analyzed the levels of PGE₂ of *G. asiatica* in June, November and February. The results showed the levels of PGE₂ were much higher in November and February than June⁽²⁾. Therefore lower temperature seemed to promote prostaglandin production. There are several factors that can be attributed to the accumulation of prostaglandins in the cold season, including stage of life cycle, activity of enzymes involved in prostaglandins synthesis, environmental conditions (seawater temperature, irradiance, contents of dissolved components) and others. Prostaglandins may be related to processes permitting the algae tissue to survive in cold water⁽³⁾. In addition, PGE₂ amount reduced when seawater temperature rose from 25°C to 35°C. The optimal growth temperature of *G. tenuistipitata* is 20-25°C. The result indicated that high temperature is not favorable for *G. tenuistipitata* and lowers the PGE₂ production.




Figure 3. PGE₂ amount of *Gracilaria tenuistipitata* in different temperatures. The seaweed was cultivated in 25°C seawater and then transferred to temperature 4°C, 15°C and 35°C for 12 hr, respectively. Values are mean ± S.D. (N = 3). a, b, c: Values with different superscripts are significantly different at $P < 0.05$.

PGE₂ amount in *G. tenuistipitata* also changed in different irradiance (Figure 4). *G. tenuistipitata* was cultivated in 330 $\mu\text{E m}^{-2} \text{s}^{-1}$ of irradiance (control condition) and then transferred to 640 $\mu\text{E m}^{-2} \text{s}^{-1}$ and dark for 12 hr. PGE₂ amount in the seaweed increased about 38% when the seaweed was transferred from 330 $\mu\text{E m}^{-2} \text{s}^{-1}$ to 640 $\mu\text{E m}^{-2} \text{s}^{-1}$ of irradiance. PGE₂ amount increased about 37% when the seaweed was transferred to dark condition. According to Imbs's report, the content of prostaglandins in *G. verrucosa* was influenced by light intensity. *G. verrucosa* grew at various levels of photosynthetic active radiation (PAR₀) including 95%, 50% and 5% PAR₀. The amount of PGE₂ and PGF_{2α} in the seaweed exposed to 50% of PAR₀ was about half of that of the control sample (95% PAR₀). But the levels of PGE₂ and PGF_{2α} were higher in extremely low (5% PAR₀) than normal illumination condition (50% PAR₀)⁽³⁾. This result was similar to that reported by Imbs *et al.* (2001).

The seaweed in this study was collected from a seaweed farm and normally fully submerged under water. PGE₂ amounts in *G. tenuistipitata* increased 25% and 31% when the seaweeds were exposed to air for 2 hr and 4 hr, respectively (Figure 5). The result was presumed that the seaweed was under stress when exposed to air. The increase of PGE₂ is responsible to adapt desiccation condition during the period of emergence.

PGE₂ amount decreased in *G. tenuistipitata* when the seaweeds were soaked into freshwater, as shown in Figure 6. PGE₂ decreased 56%, 81% and 100% when the seaweed was soaked into freshwater for 2, 4, and 8 hr, respectively. It is harmful for *G. tenuistipitata* to be soaked into freshwater. PGE₂ amount decreased gradually with increasing of the soaking time in freshwater.




Figure 4. PGE₂ amount of *Gracilaria tenuistipitata* in different irradiance. The seaweed was cultivated in seawater with irradiance 330 $\mu\text{E m}^{-2} \text{s}^{-1}$ and then transferred to 640 $\mu\text{E m}^{-2} \text{s}^{-1}$ of irradiance and dark (0 $\mu\text{E m}^{-2} \text{s}^{-1}$) for 12 hr, respectively. Values are mean ± S.D. (N = 3). a, b: Values with different superscripts are significantly different at $P < 0.05$.

The level of PGE₂ was not detected when the seaweed was soaked for 8 hr in freshwater.

Effects of metal ions on PGE₂ level of *G. tenuistipitata* were shown in Figure 7. The common amount of various metal ions in the ocean from Environmental Protection Administration, Executive Yuan, Taiwan, ROC., are 0.5, 0.03 and 0.05 mg/L (control concentration) in terms of Zn²⁺, Cu²⁺ and Mn²⁺, respectively. The concentrations of Zn²⁺, Cu²⁺ and Mn²⁺ in this study were 10 and 100-fold higher than the control concentrations. PGE₂ level did not change significantly when the concentration of Cu²⁺ in seawater was 0.3 mg/L, but was




Figure 5. PGE₂ amount of *Gracilaria tenuistipitata* in different exposure times to air. The seaweed was cultivated in seawater and then exposed to air for 2 and 4 hr, respectively. Values are mean ± S.D. (N = 3). a, b: Values with different superscripts are significantly different at $P < 0.05$.




Figure 6. PGE₂ amount of *Gracilaria tenuistipitata* in different soaking times in freshwater. The seaweed was cultivated in seawater and then soaking in freshwater for 2, 4 and 8 hr, respectively. Values are mean ± S.D. (N = 3). a, b, c: Values with different superscripts are significantly different at $P < 0.05$.

not detected when exposed into seawater with 3 mg/L of Cu²⁺ (Figure 7a). Cu²⁺ has been pointed to decrease PGE₂ synthesis. However, Cu²⁺ has no direct effect upon the transformation of AA to prostaglandin⁽¹⁵⁻¹⁷⁾. In this study, Cu²⁺ was found to block PGE₂ synthesis in *G. tenuistipitata* at higher concentration. Similarly, PGE₂ amount did not change when the concentration of Zn²⁺ in seawater was 5 mg/L, but decreased about 76% when the concentration of Zn²⁺ rose to 50 mg/L (Figure 7a). Zn²⁺ may affect prostaglandins synthesis and metabolism through regulation of enzymes related to prostaglandins synthesis⁽¹⁸⁾. In *in vitro* studies, Zn²⁺ has been reported to completely inhibit the conversion of AA to prostaglandins in peritoneal polymorphonuclear cells from rabbits^(17,19). Concentrations of Mn²⁺ had no effect on PGE₂ level (Figure 7b) as well. The common levels of Ca²⁺ and Mg²⁺ in the ocean are 400 and 1350 mg/L (control concentration), respectively. The concentration





Figure 7. Ratio of PGE₂ amount of *Gracilaria tenuistipitata* cultivated in seawater with various concentrations of metal ions. The seaweed was cultivated in seawater with various concentrations of metal ions for 12 hr. *Value is significantly different at $P < 0.05$ when compared to that of control.

of Ca^{2+} and Mg^{2+} in this study was 50% higher than the control. The result indicated that Ca^{2+} promoted PGE_2 production in *G. tenuistipitata*. PGE_2 level increased about 59% when the concentration of calcium in the seawater was 600 mg/L. We supposed that Ca^{2+} induced enzyme activity related to prostaglandins synthesis to produce more PGE_2 in *G. tenuistipitata*. However Mg^{2+} slightly decreased PGE_2 level in *G. tenuistipitata* with the concentration 2025 mg/L in the seawater (Figure 7b). The result showed that high concentration of Mg^{2+} may inhibit PGE_2 production.

G. tenuistipitata is the major species of *Gracilaria* and is commonly cultivated in Taiwan. According to the results, the environmental factors significantly affect PGE_2 production in *G. tenuistipitata*. These results can also be found in another popular seaweed (*G. verrucosa*) in Japan^(2,3), but the amount of PGE_2 was different. The PGE_2 amount in *G. verrucosa* was much higher than *G. tenuistipitata*. Furthermore, the variation of PGE_2 production in *G. verrucosa* was also greater than *G. tenuistipitata* when the seaweed grew in different environmental conditions. Therefore the PGE_2 amounts in the seaweed may also depend on species.

PGE_2 has many physiological functions which are necessary for human body, but excess level of PGE_2 is harmful. The side effects of PGE_2 included hypotension, nausea, vomiting, diarrhea etc. *G. tenuistipitata* contains low quantity of PGE_2 therefore it was not toxic in general condition. However, the environmental conditions affect the PGE_2 production in *G. tenuistipitata*, which might lead to the increase of the amount of PGE_2 in the seaweed. Hence, we suggested that minimizing the variation of growth condition could avoid PGE_2 increase in the seaweed during the cultivation period. In addition, consumers should avoid eating raw seaweed, because PGE_2 can be destroyed by heating, to reduce the amount of PGE_2 in the seaweed and may ensure the edible safety.

CONCLUSIONS

The PGE_2 production in *G. tenuistipitata* depended on environment conditions. The environmental conditions may exert effects on the PGE_2 production in the seaweed. The change in the amount of PGE_2 indicates that the seaweed adapts to different growth conditions during the period of emergence. Some special environment factors, such as low temperature, salinity, irradiance, air and Ca^{2+} , could promote PGE_2 production in the seaweed.

ACKNOWLEDGEMENTS

We appreciated to Dr. Yean-Chang Chen, Department of Aquaculture, National Taiwan Ocean University, for identifying the seaweed species. This study is supported by the grant (CMBB-97-P-B).

REFERENCES

- Noguchi, T., Matsui, T., Miyazawa, K., Asakawa, M., Iijima, N., Shida, Y., Fuse, M., Hosaka, Y., Kirigaya, C., Watabe, K., Usui, S. and Fukagawa, A. 1994. Poisoning by the red alga 'ogonori' (*Gracilaria verrucosa*) on the Nojima coast, Yokohama, Kanagawa prefecture, Japan. *Toxicol* 41: 189-194.
- Sajiki, J. 1997. Effect of acetic acid treatment on the concentrations of arachidonic acid and prostaglandin E2 in the red algae, *Gracilaria asiatica* and *G. rhodocaudata*. *Fish Sci.* 63: 128-131.
- Imbs, A. B., Vologodskaya, A. V., Nevshupova, N. V., Khotimchenko, S. V. and Titlyanoy, E. A. 2001. Response of prostaglandin content in the red alga *Gracilaria verrucosa* to season and solar irradiance. *Phytochemistry* 58: 1067-1072.
- Glickman, M. 1987. Utilisation of seaweed hydrocolloids in the food industry. *Hydrobiology* 151/152: 31-47.
- Minghetti, L. and Levi, G. 1998. Microglia as effector cells in brain damage and repair: focus on prostanooids and nitric oxide. *Prog. Neurobiol.* 54: 99-125.
- Mebane, H., Turnbach, M. E. and Randich, A. 2003. Spinal EP receptors mediating prostaglandin E2-induced mechanical hyperalgesia, thermal hyperalgesia, and touch-evoked allodynia in rats. *J. Pain* 4: 392-399.
- Lee, S. and Simon, M. D. 1999. Role and regulation of cyclooxygenase-2 during inflammation. *Am. J. Med.* 106: 37-42.
- Rocca, B. and Fitzgerald, G. A. 2002. Cyclooxygenases and prostaglandins; shaping up the immune response. *Int. Immunopharmacol.* 2: 603-630.
- Bjorkman, D. J. 1998. The effect of aspirin and nonsteroidal anti-inflammatory drug on prostaglandins. *Am. J. Med.* 105: 8-12.
- Nishihara, I., Minami, T., Uda, R., Ito, S., Hyodo, M. and Hayaishi, O. 1995. Effect of NMDA receptor antagonists on prostaglandin E2-induced hyperalgesia in conscious mice. *Brain Res.* 677: 138-144.
- Gregson, R. P., Marwood, J. F. and Quinn, R. J. 1979. The occurrence of prostaglandins PGE_2 and PGF_2 -alpha in a plant—the red alga *Gracilaria lichenoides*. *Tetraedron Lett.* 46: 4505-4506.
- Pradelles, P., Grassi, J. and Maclouf, J. A. 1985. Enzyme immunoassays of eicosanoids using acetylcholinesterase as label: an alternative to radioimmunoassay. *Anal. Chem.* 57: 1170-1173.
- Hsu, B. Y., Tsao, C. Y., Chiou, T. K. and Hwang, D. F. 2007. HPLC determination for prostaglandins from Seaweed *Gracilaria gigas*. *Food Control* 18: 639-645.
- Dawes, C. J., Kovach, C. and Friedlander, M. 1993. Exposure of *Gracilaria* to various environmental conditions. II. The effect on fatty acid composition. *Bot. Marina* 36: 289-296.
- Lands, W., Lee, R. and Smith, W. 1971. Factors regulating the biosynthesis of various prostaglandins. *Ann. N. Y. Acad. Sci.* 180: 107-122.

16. Maddox, I. S. 1973. The role of copper in prostaglandin synthesis. *Biochim. Biophys. Acta* 306: 74-81.
17. Cunnane, S. C. 1982. Differential regulation of essential fatty acid metabolism to the prostaglandins: possible basis for the interaction of zinc and copper in biological systems. *Prog. Lipid Res.* 21: 73-90.
18. Chan, J. A., Nagasawa, M., Takeguchi, C. and Sih, C. J. 1975. On agents favoring prostaglandin F formation during biosynthesis. *Biochemistry* 14: 2981-2991.
19. Mapes, C. A., Bailey, P. T., Matson, C. F., Hauer, E. C. and Sobocinski, P. Z. J. 1978. *In vitro* and *in vivo* actions of zinc ion affecting cellular substances which influence host metabolic responses to inflammation. *J. Cell. Physiol.* 95: 115-124.