



2006

## Isolation and identification of two *Antrodia cinnamomea* strains from fruiting bodies

Follow this and additional works at: <https://www.jfda-online.com/journal>

---

### Recommended Citation

Chang, C.-Y.; Lue, M.-Y.; and Pan, T.-M. (2006) "Isolation and identification of two *Antrodia cinnamomea* strains from fruiting bodies," *Journal of Food and Drug Analysis*: Vol. 14 : Iss. 2 , Article 15.  
Available at: <https://doi.org/10.38212/2224-6614.2494>

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

# Isolation and Identification of Two *Antrodia cinnamomea* Strains from Fruiting Bodies

CHIEN-YU CHANG<sup>1,2</sup>, MING-YONG LUE<sup>3</sup> AND TZU-MING PAN<sup>1\*</sup>

<sup>1</sup> Institute of Microbiology and Biochemistry, National Taiwan University, 1, Sec. 4, Roosevelt Rd., Taipei 106, Taiwan, R.O.C.

<sup>2</sup> Taiwan Sugar Research Institute, Tainan, Taiwan

<sup>3</sup> Department of Cosmetic Science, Chia-Nan University of Pharmacy and Science, Taiwan

(Received: January 26, 2005; Accepted: November 7, 2005)

## ABSTRACT

*Antrodia cinnamomea* strains ACTS1 and AC0623 were isolated from the fruiting bodies. The partial sequences of ribosomal RNA gene, including the internal transcribed spacers, ITS1-5.8S-ITS2, and 18S ribosomal RNA genes, have been sequenced and applied to identify these two strains. Four specific primer sets (NS1/NS2, NS3/NS4, NS5/NS8, and ITS1/ITS4) were utilized to perform PCR experiment of the ribosomal RNA genes. Comparison of partial nucleotide sequences (ITS1-5.8S-ITS2) with strains from different genus (*Antrodiella semisupina*, *Antrodiella romellii* and *Trametes versicolor*) was made. These DNA sequence data demonstrated that there is no difference among ACTS1, AC0623, BCRC 35396 and BCRC 35398. The colony morphology and growth characteristics of *Antrodia cinnamomea* strains in different culture media of agar plate were also made. No obvious microscopic difference of *A. cinnamomea* strains ACTS1, AC0623, BCRC 35396, and BCRC 35398 were observed. The results from this study suggested that strains ACTS1 and AC0623 isolated from fruiting bodies, and the strains BCRC 35396 and 35398, the original *A. cinnamomea* strain obtained from Food Industry Research and Development Institute (FIRDI) are the same species.

Key words: *Antrodia cinnamomea*, ribosomal RNA genes, polymerase chain reaction (PCR), fruiting bodies

## INTRODUCTION

*Antrodia cinnamomea*, a native species in Taiwan, grows naturally on *Cinnamomum kanehirai* Hay, a kind of camphor tree grown only in Taiwan, and is commonly known as "Niu-chang-chih". This fungus forms fruit bodies that causes brown rot on *C. kanehirai*<sup>(1,2,3)</sup>. Aborigines living in Taiwan's mountain range have used this fungus for the treatment of food and drug intoxications, diarrhea, abnormal pains, hypertension and liver cancer in Taiwanese folk medicine<sup>(2,4,5,6)</sup>. The current price of fresh fruiting bodies of *A. cinnamomea* is estimated to be US\$ 1500 per kg.

Economic incentives are effective for the development of artificial cultivation methods. Some organizations claim to have induced formation of fruiting bodies since early 2001. However, it seems that growth under such artificial environments still depends on luck. Nevertheless, submerged mycelia products have been recently marketed. We have isolated two strains ACTS1 and AC0623 from fruiting bodies of *A. cinnamomea* at Liou Kuei in southern Taiwan. In order to understand whether strains ACTS1 and AC0623 cultivated in Taiwan Sugar Research Institute (TSRI) and the BCRC 35396 and 35398 strains are the same, some molecular systematic techniques were made<sup>(7-22)</sup>. Random amplified polymorphic DNA (RAPD) markers with polymerase chain reaction (PCR)

amplification of genomic DNA technique were also used for identification of *Cordyceps* species<sup>(17,23,24)</sup>.

Based on PCR fragments of the whole ribosomal RNA gene (including the internal transcribed spacers [ITS1 and ITS2], plus 5.8S, 18S, and 28S ribosomal RNA genes) we can distinguish whether these strains have the same characteristics from the original species. The results of isolation and identification of these strains are described as follows.

## MATERIALS AND METHODS

### I. Strains of *Antrodia cinnamomea*

The strains of this study *A. cinnamomea* BCRC 35396 and *A. cinnamomea* BCRC 35398 were purchased from Bioresources Collection & Research Center (BCRC, FIRDI, Hsinchu, Taiwan). Strains of ACTS1 and AC0623 were isolated from fruiting bodies of *A. cinnamomea* at Liou Kuei in southern Taiwan as illustrated in Table 1.

### II. Isolation of Two *Antrodia cinnamomea* Strains from Fruiting Bodies

Firstly the fruiting body was treated with antibiotics penicillin V or ampicillin to inhibit the growth of contaminants. Then it was cut into small pieces, put into the sterilized water and isolated from different dilutions of sterilized water on agar plate. Colonies were grown

\* Author for correspondence. Tel: +886-2-33664519; Fax: +886-2-2362-7044; E-mail: tmpan@ntu.edu.tw

on plate for three weeks on M25 plate. Two strains of ACTS1 and AC0623 were isolated from fruiting bodies of *A. cinnamomea*.

### III. *Antrodia cinnamomea* Cultured Media

*A. cinnamomea* strains ACTS1, AC0623, BCRC 35396, and BCRC 35398 were cultured and maintained at 28°C on M25 agar plate (Merck, Germany). Mycelium obtained from the edge of 21-day-old colonies was used as the inoculum source. Other culture media such as PDA (potato dextrose agar, Difco, Becton Dickinson, USA), OAT (oatmeal agar, Difco, Becton Dickinson, USA), SGA (Sabouraud-2% glucose agar, Merck, Germany), and NA (nutrient agar, Merck, Germany) agar plate were used to make a comparison of colony growth. Cell mass production was carried out in 500-mL Hinton flask with rotary shaker of 110 rpm and 28°C respectively. The M25 cultured media of *A. cinnamomea* strains are as follows: glucose 3%, bacto-peptone 1.5%, yeast extract 1.5%, malt extract 1.5%, MgSO<sub>4</sub>·7H<sub>2</sub>O 0.03%, KH<sub>2</sub>PO<sub>4</sub> 0.03% and K<sub>2</sub>HPO<sub>4</sub> 0.03%, pH 5.5.

### IV. Extraction of Genomic DNA

Genomic DNA was prepared using plant DNAzol Reagent<sup>(25)</sup> (Invitrogen, USA). The procedures were as follows.

#### (I) Extration

Plant tissues of *A. cinnamomea* were pulverized in liquid nitrogen using mortar and pestle. As the samples were grinded, liquid nitrogen was replenished in the mortar 2 to 3 times. The resulting homogenous frozen powder was then transferred to a microcentrifuge tube containing plant DNAzol. Every gram of plant tissue needed 3 mL of plant DNAzol. The mixture was mixed thoroughly by gentle inversion a few times before incubation at 25°C and shaking for 5 min. The mixture was then added to 0.3 mL of chloroform and mixed vigorously. After incubation at 25°C for another 5 min, the mixture was centrifuged.

#### (II) Centrifugation

Following extraction, the extract was centrifuged at 12,000 × g for 10 min. The supernatant was transfer to a fresh tube.

#### (III) DNA Precipitation

The collected supernatant was mixed with 0.23 mL of absolute ethanol. The compound was mixed by inverting the tube 6 to 8 times before storage at room temperature for 5 min. Precipitated DNA was separated by centrifugation at 5,000 × g for 4 min.

**Table 1.** Source and collection number of *Antrodia cinnamomea* strains in this study

Species	Collection number	Source
<i>Antrodia cinnamomea</i>	BCRC 35396	BCRC <sup>a</sup> , Taiwan
<i>Antrodia cinnamomea</i>	BCRC 35398	BCRC, Taiwan
<i>Antrodia cinnamomea</i> fruiting body	ACTS1	Liou Kuei, Taiwan isolated by TSRI <sup>b</sup> , Taiwan
<i>Antrodia cinnamomea</i> fruiting body	AC0623	Liou Kuei, Taiwan, isolated by TSRI, Taiwan

<sup>a</sup>BCRC: Bioresources Collection & Research Center, Hsinchu, Taiwan.

<sup>b</sup>TSRI: Taiwan Sugar Research Institute, Tainan, Taiwan.

**Table 2.** Primers used in this study<sup>(21)</sup>

Primer designation	Primer sequences (5'→3')	Tm (°C)
NS1	GTAGTCATATGGCTTGTCTC	54
NS2	GGCTGCTGGCACCAGACTTGC	62
NS3	GCAAGTCTGGTGCCAGCAGCC	62
NS4	CTCCCGTCAATTCCTTTAAG	56
NS5	AACTTAAAGGAATTGACGGAAG	49
NS8	TCCGCAGGTTACCTACGGA	64
ITS1	TCCGTAGGTGAACCTGCGG	62
ITS4	TCCTCCGCTTATTGATATGC	58

#### (IV) DNA Wash

##### 1. Plant DNAzol-ethanol Wash

Plant DNAzol-ethanol wash was prepared by mixing 1 volume of DNAzol with 0.75 volume of 100% ethanol. A quantity of 0.3 mL of Plant DNAzol-ethanol wash solution was mixed with the DNA precipitate by vortexing. After 5 min, the mixture was subjected to centrifugation at 5,000 × g for 4 min.


##### 2. Ethanol Wash

After the DNAzol wash solution was removed, the DNA pellet was washed by vigorously mixing with 0.3 mL of 75% ethanol. The mixture was then centrifuged at 5,000 × g for 4 min.

#### (V) DNA Solubilization

After ethanol layer was removed by decanting, the remaining ethanol was further removed with a micropipette that was allowed to sit vertically for about 2 min. The remaining DNA pellet was air dried and then dissolved in 70 µL of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0). If DNA pellet is difficult to dissolve, 8 mM NaOH solution can be used instead of TE buffer. In a typical DNA preparation, the DNA solution is cloudy and





**Figure 3.** Anatomy microscopic characteristics of *Antrodia cinnamomea* (A) ACTS1 (B) AC0623 (C) BCRC 35396 (D) BCRC 35398 on M25 agar ( $\times 50$ ).

**Table 3.** Diameter of the colonies of *Antrodia cinnamomea* strains in different media

Strain	ACTS1	AC0623	BCRC 35396	BCRC 35398
Medium				
M25	5.0 cm <sup>a</sup>	6.5 cm	6.0 cm	5.0 cm
PDA	4.5 cm	5.5 cm	5.5 cm	5.3 cm
OAT	4.5 cm	4.5 cm	5.3 cm	6.5 cm
SGA	4.3 cm	4.5 cm	4.0 cm	5.0 cm
NA	— <sup>b</sup>	—	—	—

<sup>a</sup>cm: the length were measured when the colonies reached their maximum diameter of *A. cinnamomea* strains within 21 days after inoculated on the plates.

<sup>b</sup>No growth.

The growth rate of *A. cinnamomea* on M25 agar was the fastest among all combination tested. The microscopic traits of these strains are shown in Figure 2, and the anatomy microscopic characteristics of *A. cinnamomea* are depicted in Figure 3.


## II. Comparison of the PCR Products of Ribosomal RNA Genes among the *Antrodia cinnamomea* Strain ACTS1, AC0623, BCRC 35396, and BCRC 35398

Four specific primer sets (NS1/NS2, NS3/NS4, NS5/NS8, and ITS1/ITS4) were utilized to perform PCR amplification of the ribosomal RNA genes (including 18S and ITS1-5.8S-ITS2 rRNA gene). The results are shown in Figure 4. The fragment length and location with specific primer sets to perform PCR amplification are shown as follows:

The PCR product of NS1 & NS2: 550 bp (18S rRNA gene)

The PCR product of NS3 & NS4: 600 bp (18S rRNA gene)

The PCR product of NS5 & NS8: 650 bp (18S rRNA gene)



**Figure 4.** Comparison of the PCR products of ribosomal RNA genes among the *Antrodia cinnamomea* strain ACTS1, AC0623 (TSRI), and BCRC 35396, BCRC 35398 strain (FIRDI). PCR products were separated by electrophoresis in a 2.0% agarose gel and then stained with ethidium bromide. The gel was observed and photographed under ultraviolet light. The 100-bp ladder (M) as molecular weight standards is shown in intervals.

The PCR product of ITS1 & ITS4: 700 bp (ITS1-5.8S-ITS2 rRNA gene)

## III. DNA Sequence Analysis of *Antrodia cinnamomea* Strain<sup>(16)</sup> and Others

Four specific primer sets (NS1/NS2, NS3/NS4, NS5/



NS8, and ITS1/ITS4) were utilized to perform primer for direct DNA sequence of PCR product. Ribosomal RNA genes of ACTS1, BCRC 35396, BCRC 35398, and

AC0623 were used for DNA sequence analysis. Based on the ribosomal RNA genes (18S and ITS1-5.8S-ITS2 rRNA genes) and four specific primer sets, the length of PCR product after reaction was 556 bp, 602 bp, 659 bp and 682 bp respectively. The complete sequence of 18S rRNA, ITS1, 5.8S rRNA and ITS2 were obtained. The results of the DNA sequences analysis among ACTS1, AC0623, BCRC 35396, and BCRC 35398 strains are shown in Table 4.

**Table 4.** Similarity of DNA sequences of partial ribosomal RNA genes [containing partial 18S ribosomal RNA gene, internal transcribed spacer 1 (ITS1), 5.8S ribosomal RNA gene, internal transcribed spacer 2 (ITS2), and partial 28S ribosomal RNA gene] among 4 *Antrodia cinnamomea* strains including ACTS1 (AY378092), AC0623 (AY378093), BCRC 35396 (AY378094), and BCRC 35398 (AY378095)

Strain	AC0623	BCRC 35396	BCRC 35398	ACTS1
ACTS1	100%	99.8%	99.9%	100%
AC0623	100%	99.8%	99.9%	100%
BCRC 35396	99.8%	100%	99.9%	99.8%
BCRC 35398	99.9%	99.9%	100%	99.9%

From the sequence analysis of *Antrodia cinnamomea*<sup>(26)</sup>, 18S rRNA is shown to be a high conservative region (about 1700 bp) and the similarity between BCRC 35396 and BCRC 35398 strains was over 98%. Since the region of partial nucleotide sequence ITS1-5.8S-ITS2 (about 680 bp) is non-conservative evolutionally, it is suitable for species identification. Partial nucleotide sequences (ITS1-5.8S-ITS2) were compared with those of the strains from different genus (*Antrodiella semisupina*, *Antrodiella romellii*, and *Trametes versicolor*). The sequences were analyzed

BCRC35396	TCCGTAGGTGAACCTGCGGAAGGACCATTATTGTAT---TTGAAAGGGG
BCRC35716	TCCGTAGGTGAACCTGCGGAAGGACCATTATTGTAT---TTGAAAGGGG
AC0623	TCCGTAGGTGAACCTGCGGAAGGATCATTATTGTAT---TTGAAAGGGG
ACTS1	TCCGTAGGTGAACCTGCGGAAGGATCATTATTGTAT---TTGAAAGGGG
BCRC35398	TCCGTAGGTGAACCTGCGGAAGGATCATTATTGTAT---TTGAAAGGGG
<i>Antrodiella semisupina</i>	TCCGTAGGTGAACCTGCGGAAGGATCATTAAATGAATGAACTTGGGCATAG
<i>Antrodiella romellii</i>	TCCGTAGGTGAACCTGCGGAAGGATCATTAAATGAATGAACTTGGGCAAAG
<i>Trametes versicolor</i>	TCCGTAGGTGAACCTGCGGAAGGATCATTAAACGAGT---TTTAAAACGAG
BCRC35396	TTGTAGCTGACCTCCTCTTGAAAAGGGGGGAGGTATGTGCACACCTCT-G
BCRC35716	TTGTAGCTGACCTCCTCTTGAAAAGGGGGGAGGTATGTGCACACCTCTTG
AC0623	TTGTAGCTGACCTCCTCTTGAAAAGGGGGGAGGTATGTGCACACCTCT-G
ACTS1	TTGTAGCTGACCTCCTCTTGAAAAGGGGGGAGGTATGTGCACACCTCT-G
BCRC35398	TTGTAGCTGACCTCCTCTTGAAAAGGGGGGAGGTATGTGCACACCTCT-G
<i>Antrodiella semisupina</i>	TTGTAGCTGGCCTC-----AGCAATGAGGCATGTGCACAC-TGT-G
<i>Antrodiella romellii</i>	TTGTAGCTGGCCTC-----AGCAATGGGGCATGTGCACAC-TTT-G
<i>Trametes versicolor</i>	TTGTAGCTGGCCTT-----C---CGGGGCATGTGCACGC-TCT-G
BCRC35396	TTCATTCATATTCTCTCACACCTGTGCATGCTTTGTAGGTTGGTTTTGAA
BCRC35716	TTCATTCATATTCTCTCACACCTGTGCATGCTTTGTAGGTTGGTTTTGAA
AC0623	TTCATTCATATTCTCTCACACCTGTGCATGCTTTGTAGGTTGGTTTTGAA
ACTS1	TTCATTCATATTCTCTCACACCTGTGCATGCTTTGTAGGTTGGTTTTGAA
BCRC35398	TTCATTCATATTCTCTCACACCTGTGCATGCTTTGTAGGTTGGTTTTGAA
<i>Antrodiella semisupina</i>	TTCATCCAC-CCTTCACACCTCTGTGCACTTCTCATGGGTTTGGGTCAAG
<i>Antrodiella romellii</i>	TTCATCCAC-C-TTCACACCACTGTGCACTTCTCATGGGTC-GGGTTGCG
<i>Trametes versicolor</i>	CTCATCCAC-TCT---ACCCCTGTGCACTTACTGTAGGTTGGCGTGGGC
BCRC35396	TGGTTGTCTTCTCTGATGGAGACAGCTGTTTTGACCTTCCTATGTTTTTT
BCRC35716	TGGTTGTCTTCTCTGATGGAGACAGCTGTTTTGACCTTCCTATGTTTTTT
AC0623	TGGTTGTCTTCTCTGATGGAGACAGCTGTTTTGACCTTCCTATGTTTTTT
ACTS1	TGGTTGTCTTCTCTGATGGAGACAGCTGTTTTGACCTTCCTATGTTTTTT
BCRC35398	TGGTTGTCTTCTCTGATGGAGACAGCTGTTTTGACCTTCCTATGTTTTTT
<i>Antrodiella semisupina</i>	T-----CTGAAATATGGCGAAGCCC-----CCTTCTCATGTGTTTT
<i>Antrodiella romellii</i>	T-----CTGAAATATGGCAAAGCCC-----TTCTCATGTGTTTT
<i>Trametes versicolor</i>	T-----CCTAACGGGAGCATTCTG-----CCGGCCTATGTATACT

BCRC35396	AAATTGACTCCGTATCAGTTACAGAATGTA----TGTTGCGTGTAACGCA
BCRC35716	AAATTGACTCCGTATCAGTTACAGAATGTA----TGTTGCGTGTAACGCA
AC0623	AAATTGACTCCGTATCAGTTACAGAATGTA----TGTTGCGTGTAACGCA
ACTS1	AAATTGACTCCGTATCAGTTACAGAATGTA----TGTTGCGTGTAACGCA
BCRC35398	AAATTGACTCCGTATCAGTTACAGAATGTA----TGTTGCGTGTAACGCA
<i>Antrodiella semisupina</i>	AAACACACACTATAACAAGTTTTAGAAATGTAACAATCATGCGTC-AATGCA
<i>Antrodiella romellii</i>	AC-CACACACT--ACAAGTTTTAGAAATGTAACAATCATGCATT-AATGCA
<i>Trametes versicolor</i>	ACAAACACTTTA---AAGTATCAGAATGTAA-----ACGCGTCTAACGCA
BCRC35396	T--ATTGTATAACTTTTCAGCAACGGATCTCTTGGCTCTCGCATCGATGAA
BCRC35716	T--ATTGTATAACTTTTCAGCAACGGATCTCTTGGCTCTCGCATCGATGAA
AC0623	T--ATTGTATAACTTTTCAGCAACGGATCTCTTGGCTCTCGCATCGATGAA
ACTS1	T--ATTGTATAACTTTTCAGCAACGGATCTCTTGGCTCTCGCATCGATGAA
BCRC35398	T--ATTGTATAACTTTTCAGCAACGGATCTCTTGGCTCTCGCATCGATGAA
<i>Antrodiella semisupina</i>	T--ATAATACAACCTTTTCAGCAACGGATCTCTTGGCTCTCGCATCGATGAA
<i>Antrodiella romellii</i>	T--ATAATACAACCTTTTCAGCAACGGATCTCTTGGCTCTCGCATCGATGAA
<i>Trametes versicolor</i>	TCTATAACAACCTTTTAGCAACGGATCTCTTGGCTCTCGCATCGATGAA
BCRC35396	GAACACGGCGAAATGTGATAAGTAATGTGAATTGCAGAATTCAGTGAATC
BCRC35716	GAACACGGCGAAATGTGATAAGTAATGTGAATTGCAGAATTCAGTGAATC
AC0623	GAACACGGCGAAATGTGATAAGTAATGTGAATTGCAGAATTCAGTGAATC
ACTS1	GAACACGGCGAAATGTGATAAGTAATGTGAATTGCAGAATTCAGTGAATC
BCRC35398	GAACACGGCGAAATGTGATAAGTAATGTGAATTGCAGAATTCAGTGAATC
<i>Antrodiella semisupina</i>	GAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTCAGTGAATC
<i>Antrodiella romellii</i>	GAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTCAGTGAATC
<i>Trametes versicolor</i>	GAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTCAGTGAATC
BCRC35396	ATCGAATCTTTGAACGCACCTTGCGCTCCTTGGTATTCTGAGGAGCATGC
BCRC35716	ATCGAATCTTTGAACGCACCTTGCGCTCCTTGGTATTCTGAGGAGCATGC
AC0623	ATTGAATCTTTGAACGCACCTTGCGCTCCTTGGTATTCTGAGGAGCATGC
ACTS1	ATTGAATCTTTGAACGCACCTTGCGCTCCTTGGTATTCTGAGGAGCATGC
BCRC35398	ATCGAATCTTTGAACGCACCTTGCGCTCCTTGGTATTCTGAGGAGCATGC
<i>Antrodiella semisupina</i>	ATCGAATCTTTGAACGCACCTTGCGCTCCTTGGTATTCCGAGGAGCATGC
<i>Antrodiella romellii</i>	ATCGAATCTTTGAACGCACCTTGCGCTCCTTGGTATTCCGAGGAGCATGC
<i>Trametes versicolor</i>	ATCGAATCTTTGAACGCACCTTGCGCTCCTTGGTATTCCGAGGAGCATGC
BCRC35396	CTGTTTGAGTGTCATGGAATTATCAACCCTTTTGACTTTTTTGT-----
BCRC35716	CTGTTTGAGTGTCATGGAATTATCAACCCTTTTGACTTTTTTGT-----
AC0623	CTGTTTGAGTGTCATGGAATTATCAACCCTTTTGACTTTTTTGT-----
ACTS1	CTGTTTGAGTGTCATGGAATTATCAACCCTTTTGACTTTTTTGT-----
BCRC35398	CTGTTTGAGTGTCATGGAATTATCAACCCTTTTGACTTTTTTGT-----
<i>Antrodiella semisupina</i>	CTGTTTGAGTGTCATGGTATTCTCAACCCTGCTACATTTTTTTGAAAGAT
<i>Antrodiella romellii</i>	CTGTTTGAGTGTCATGGTATTCTCAACCCTGCTACATTTTTTT-----
<i>Trametes versicolor</i>	CTGTTTGAGTGTCATGGAATTCTCAACTT--ATAAATCCTTGTG-----
BCRC35396	-TGAATGGGCTTGGATTTGGAGGGTTAAATTGCTGGCTCTT---CTTTT
BCRC35716	-TGAATGGGCTTGGATTTGGAGGGTTAAATTGCTGGCTCTT---TTTTT
AC0623	-TGAATGGGCTTGGATTTGGAGGGTTAAATTGCTGGCTCTT---TTTTT
ACTS1	-TGAATGGGCTTGGATTTGGAGGGTTAAATTGCTGGCTCTT---TTTTT
BCRC35398	-TGAATGGGCTTGGATTTGGAGGGTTAAATTGCTGGCTCTT---TTTTT
<i>Antrodiella semisupina</i>	GTTGCTTGGCTTGGACTTGGAGGTTT--ATTGCTGGCATTCAACTTGTTT
<i>Antrodiella romellii</i>	GTAGCTGGGCTTGGACTTGGAGGT---ATTGCCGGTGTTC---TCTTTT
<i>Trametes versicolor</i>	ATCTATAAGCTTGGACTTGGAGGC-----TTGCTGGCCCTT-----

BCRC35396	T-GATTCAGCTCCTCTTGAATGCATTAGCTTGAACCCTTGTGGATTGACC
BCRC35716	T-GATTCAGCTCCTCTTGAATGCATTAGCTTGAACCCTTGTGGATTGACC
AC0623	TTGATTCAGCTCCTCTTGAATGCATTAGCTTGAACCCTTGTGGATTGACC
ACTS1	TTGATTCAGCTCCTCTTGAATGCATTAGCTTGAACCCTTGTGGATTGACC
BCRC35398	T-GATTCAGCTCCTCTTGAATGCATTAGCTTGAACCCTTGTGGATTGACC
<i>Antrodiella semisupina</i>	GAACGCGGGCTCCTCTGAAATGCATTAGCTGGAATGT-TACCGAGCATGA
<i>Antrodiella romellii</i>	GAACGCGGGCTCCTCTGAAATGCATTAGCTGGAATGT-TACCGAGCATGA
<i>Trametes versicolor</i>	-GCGGTCCGGCTCCTCTTGAATGCATTAGCTTGAATCCGTACGGATCGGCT
BCRC35396	TTATCGGTGTGATAGTCATCTATGCCGTGGTGTCTGAGGTGGGATCGGC
BCRC35716	TTATCGGTGTGATAGTCATCTATGCCGTGGTGTCTGAGGTGGGATCGGC
AC0623	TTATCGGTGTGATAGTCATCTATGCCGTGGTGTCTGAGGTGGGATCGGC
ACTS1	TTATCGGTGTGATAGTCATCTATGCCGTGGTGTCTGAGGTGGGATCGGC
BCRC35398	TTATCGGTGTGATAGTCATCTATGCCGTGGTGTCTGAGGTGGGATCGGC
<i>Antrodiella semisupina</i>	TT--CAATGTGATAATTGTCTACGTTGCTTCAACTTGGTATTAATTCTGT
<i>Antrodiella romellii</i>	TT--CAATGTGATAATTGTCTACGTTGCTTCAACTTGGTATTAATTCTGT
<i>Trametes versicolor</i>	CT--CAGTGTGATAATTGTCTACGCTGT--GACCGTGAAGTGTTTTGGC
BCRC35396	TTCTAATGGTGCAAGTCCCTTCAGGGGGATGATTTTCTAATGACCTTCTG
BCRC35716	TTCTAATGGTGCAAGTCCCTTCAGGGGGATGATTTTCTAATGACCTTCTG
AC0623	TTCTAATGGTGCAAGTCCCTTCAGGGGGATGATTTTCTAATGACCTTCTG
ACTS1	TTCTAATGGTGCAAGTCCCTTCAGGGGGATGATTTTCTAATGACCTTCTG
BCRC35398	TTCTAATGGTGCAAGTCCCTTCAGGGGGATGATTTTCTAATGACCTTCTG
<i>Antrodiella semisupina</i>	TTCAGCTTCTAACCGTCCCTTTGCGGGACAATATCT---TGAACATCTG
<i>Antrodiella romellii</i>	TTCAGCTTCTAACCGTCCGCA---AGGACAATATCT---TGAACATCTG
<i>Trametes versicolor</i>	--GAGCTTCTAACCGTCCATT---AGGACAATTCT---T-AACATCTG
BCRC35396	ACCTCAAATCAGGTAGGACTACCCGCTGAACTTAAGCATATCAATAAGCG
BCRC35716	ACCTCAAATCAGGTAGGACTACCCGCTGAACTTAAGCATATCAATAAGCG
AC0623	ACCTCAAATCAGGTAGGACTACCCGCTGAACTTAAGCATATCAATAAGCG
ACTS1	ACCTCAAATCAGGTAGGACTACCCGCTGAACTTAAGCATATCAATAAGCG
BCRC35398	ACCTCAAATCAGGTAGGACTACCCGCTGAACTTAAGCATATCAATAAGCG
<i>Antrodiella semisupina</i>	ACCTCAAATCAGGTAGGACTACCCGCTGAACTTAAGCATATCAATAAGCG
<i>Antrodiella romellii</i>	ACCTCAAATCAGGTAGGACTACCCGCTGAACTTAAGCATATCAATAAGCG
<i>Trametes versicolor</i>	ACCTCAAATCAGGTAGGACTACCCGCTGAACTTAAGCATATCAATAAGCG
BCRC35396	GAGGA
BCRC35716	GAGG-
AC0623	GAGGA
ACTS1	GAGGA
BCRC35398	GAGGA
<i>Antrodiella semisupina</i>	GAGGA
<i>Antrodiella romellii</i>	GAGGA
<i>Trametes versicolor</i>	GAGGA

**Figure 5.** Comparison of DNA sequences of partial ribosomal RNA genes (containing partial ITS1, 5.8S rRNA gene, and partial ITS2) among 5 *Antrodia cinnamomea* strains including ACTS1 (AY378092), AC0623 (AY378093), BCRC 35396 (AY378094), BCRC 35398 (AY378095), and BCRC 35716 (AJ496402), as well as *Antrodiella semisupina* (AF126903), *Antrodiella romellii* (AF126902), and *Trametes versicolor* (AY309015) strains.

and are shown in Figure 5. The results from this study suggest that the two strains (ACTS1 and AC0623) isolated from fruiting bodies are of the same species, *A. cinnamomea*. The nucleotide sequences reported in this and other paper have been submitted to the GenBank with accession numbers AY378092 (ACTS1), AY378093

(AC0623), AY378094 (BCRC 35396), AY378095 (BCRC 35398), AJ496402 (BCRC 35716)<sup>(26)</sup>, AY309015 (*Trametes versicolor*), AF126902 (*Antrodiella romellii*)<sup>(27)</sup>, and AF126903 (*Antrodiella semisupina*)<sup>(27)</sup>.



## DISCUSSION

PCR primers that target conserved regions of fungal rRNA genes, amplifying sequence-variable fragments of genes or intervening noncoding regions, have been used for sequence comparisons. Interspecies variability is also manifested in the fragment size of the internal transcribed spacer 1 and 2 (ITS1 and ITS2, respectively) regions<sup>(28)</sup>. In order to identify *A. cinnamomea* rapidly and correctly, the molecular biology methods for *A. cinnamomea* species and strains analysis were used<sup>(26)</sup>. Firstly, they used 18S rRNA gene, ITS gene and intergenic spacer of rRNA, and Mn-superoxide dismutase gene<sup>(26)</sup>. In addition, phylogenetic analysis methods, Neighbor Joining method, Parsimony method and Maximum Likelihood method were also used to analyze evolution relation of *A. cinnamomea*<sup>(26)</sup>. The specific primer sets based PCR fragments of the whole ribosomal RNA gene [including the internal transcribed spacers (ITS1 and ITS2), 5.8S, and 18S ribosomal RNA genes] combined partial nucleotide sequence (ITS1-5.8S-ITS2) analysis with other genus for negative control could be applied to investigate whether the strain cultivated in Taiwan Sugar Research Institute (TSRI) is the species named *A. cinnamomea*. Comparison of colony morphology and growth characteristics of *Antrodia cinnamomea* strains in different culture media of agar plates was made. The growth rate of *A. cinnamomea* on M25 agar was the fastest among all combinations tested. Nutrient agar medium was not suitable for *A. cinnamomea* strains. The results demonstrate that there is no difference in the electrophoresisgram of the amplified PCR products between strains ACTS1, AC0623, BCRC 35396, and BCRC 35398. No obvious microscopic difference of *A. cinnamomea* strains ACTS1, AC0623, BCRC 35396, and BCRC 35398 was observed in this study. The major biologically active component triterpenoids will be in conjunction with our continuing research from the cultured *A. cinnamomea* mycelia.

## ACKNOWLEDGEMENTS

We would like to thank Mr. Shun-Yuan Yang for assisting the PCR experiment in this study.

## REFERENCES

- Andersson, B. E., Tornberg, K., Henrysson, T. and Olsson, S. 2001. Three-dimensional outgrowth of a wood-rotting fungus added to a contaminated soil from a former gasworks site. *Bioresour. Technol.* 78: 37-45.
- Chang, T. T. and Chou, W. N. 1995. *Antrodia cinnamomea* sp. nov. on *Cinnamomea kanehirai* in Taiwan. *Mycol. Res.* 99: 756-758.
- Wu, S. H., Ryvarden, L. and Chang, T. T. 1997. *Antrodia cinnamomea* ("niu-chang-chih"), new combination of a medicinal fungus in Taiwan. *Bot. Bull. Acad. Sin.* 38: 273-275.
- Cherng, I. H. and Chiang, H. C. 1995. Three new triterpenoids from *Antrodia cinnamomea*. *J. Natural. Products* 58: 365-371.
- Hseu, Y. C., Chang, W. C., Hseu, Y. T., Lee, C. Y., Tech, Y. J., Chen, P. C., Chen, J. Y. and Yang, H. L. 2002. Protection of oxidative damage by aqueous extract from *Antrodia cinnamomea* mycelia in normal human erythrocytes. *Life Sci.* 71: 469-482.
- Song, T. Y. and Yen, G. C. 2002. Antioxidant properties of *Antrodia cinnamomea* in submerged culture. *J. Agric. Food Chem.* 50: 3322-3327.
- Buzina, W., Doris, L. L., Braun, H., Freudenschuss, K. and Stammberger, H. 2001. Development of molecular methods for identification of *Schizophyllum commune* from clinical samples. *J. Clin. Microbiol.* 39: 2391-2396.
- Chen, Y. Q., Wang, N., Qu, L. H., Li, T. H. and Zhang, W. M. 2001. Determination of the anamorph of *Cordyceps sinensis* inferred from the analysis of the ribosomal DNA internal transcribed spacers and ITS1-5.8S-ITS2 rDNA. *Biochem. Syst. Ecol.* 29: 597-607.
- Chen, Y. Q., Wang, N., Zhou, H. and Qu, L. H. 2002. Differentiation of medicinal *Cordyceps* species by rDNA ITS sequence analysis. *Planta. Med.* 68: 635-639.
- Eisenga, B. H., Trestrail III, J. H. and Emerson, T. S. 1998. Identification of unknown mushrooms: the good, the bad, and the ugly. *J. Toxicol. Clin. Toxicol.* 36: 635-636.
- Henry, T., Iwen, P. C. and Hinrichs, S. H. 2000. Identification of *Aspergillus* species using internal transcribed spacer regions 1 and 2. *J. Clin. Microbiol.* 38: 1510-1515.
- Jackson, C. J. 2001. Molecular identification and strain typing of dermatophyte fungi. *Nippon Ishinkin Gakkai Zasshi* 42: 7-10.
- James, T. Y., Moncalvo, J. M., Li, S. and Vilgalys, R. 2001. Polymorphism at the ribosomal DNA spacers and its relation to breeding structure of the widespread mushroom *Schizophyllum commune*. *Genetics* 157: 149-161.
- Kelly, J. M. and Cox, R. A. 1982. The nucleotide sequence at the 3'-end of *Neurospora crassa* 18S-rRNA and studies on the interaction with 5S-rRNA. *Nucleic Acids Res.* 10: 6733-6745.
- Lee, S. B. and Taylor, J. W. 1990. Isolation of DNA from fungal mycelia and single spores. In "PCR Protocols: A Guide to Methods and Applications". 2nd ed. pp. 315-322. Academic Press. San Diego, U. S. A.
- Lott, T. J., Burns, B. M., Zancoppe-Oliveira, R., Elie, C. M. and Reiss, E. 1998. Sequence analysis of the internal transcribed spacer 2 (ITS2) from yeast species within the genus *Candida*. *Curr. Microbiol.* 36: 63-69.
- Lue, M. Y., Chang, C. C. and Liou, J. J. 2002. Identification of *Cordyceps* species. *Rept. Taiwan Sugar Res. Inst.* 175-176: 51-58.

18. Turenne, C. Y., Sanche, S. E., Hoban, D. J., Karlowsky, J. A. and Kabani, A. M. 1999. Rapid identification of fungi by using the ITS2 genetic region and an automated fluorescent capillary electrophoresis system. *J. Clin. Microbiol.* 37: 1846-1851.
19. Varga, J., Toth, B., Rigo, K., Teren, J., Hoekstra, R. F. and Kozakiewicz, Z. 2000. Phylogenetic analysis of *Aspergillus* section *circumdati* based on sequences of the internal transcribed spacer regions and the ITS1-5.8S-ITS2 rRNA gene. *Fungal Genet. Biol.* 30: 71-80.
20. Velegraki, A., Kambouris, M., Kostourou, A., Chalevelakis, G. and Legakis, N. J. 1999. Rapid extraction of fungal DNA from clinical samples for PCR amplification. *Med. Mycol.* 37: 69-73.
21. Vilgalys, R. and Hester, M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *J. Bacteriol.* 172: 4238-4246.
22. Zhang, D., Yang, Y., Castlebury, L. A. and Cerniglia, C. E. 1996. A method for the large scale isolation of high transformation efficiency fungal genomic DNA. *FEMS. Microbiol. Lett.* 145: 261-265.
23. Chen, Y., Zhang, Y. P., Yang, Y. and Yang, D. 1999. Genetic diversity and taxonomic implication of *Cordyceps sinensis* as revealed by RAPD markers. *Biochem. Genet.* 37: 210-213.
24. Chen, Y., Wang, W., Yang, Y. B., Su, Y. P., Zhang, L., Xiong, Z., He, C. S. and Yang, D. 1997. Genetic divergence of *Cordyceps sinensis* as estimated by random amplified polymorphic DNA analysis. *Acta. Genet. Sinica.* 24: 410-416.
25. Chomczynski, P., Mackey, K., Drews, R. and Wilfinger, W. 1997. DNAzol: a reagent for the rapid isolation of genomic DNA. *Biotechniques* 22: 550-553.
26. Chang, Y. H. 2002. Molecular systematics of *Antrodia cinnamomea*. Master Thesis, Graduate Institute of Agricultural Chemistry, National Taiwan University, Taiwan, R. O. C.
27. Johannesson, H., Renvall, P. and Stenlid, J. 2000. Taxonomy of *Antrodiella* inferred from morphological and molecular data. *Mycol. Res.* 104(1): 92-99.
28. Williams, D. W., Wilson, M. J., Lewis, M. A. O. and Potts, A. J. C. 1995. Identification of *Candida* species by PCR and restriction fragment length polymorphism analysis of intergenic spacer regions of ribosomal DNA. *J. Clin. Microbiol.* 33: 2476-2479.